

ON A DEFORMED TOPOLOGICAL VERTEX

JIAN ZHOU

ABSTRACT. We introduce a deformed topological vertex and use it to define deformations of the topological string partition functions of some local Calabi-Yau geometries. We also work out some examples for which such deformations satisfy a deformed Gopakumar-Vafa integrality and can be identified with the equivariant indices of some naturally defined bundles on the framed moduli spaces.

1. INTRODUCTION

Since the introduction of the topological vertex in [1], there have been interests in generalizing it to have more variables. Very recently one such generalization using Macdonald's polynomials has been introduced [4]. In this paper we will consider another one.

Recall the topological vertex involves three partitions, when one of them is empty it is related to the leading term of the large N colored HOMFLY polynomials of the Hopf link, and the general topological vertex can be expressed in terms of such leading terms. We will use the whole colored HOMFLY polynomials to define the deformed vertex, hence deformations of the partition functions of topological strings.

We define the deformed topological string partition functions simply by replacing the topological vertices by the deformed vertices, so their geometric meaning is missing at present. Nevertheless, they share some important properties with the undeformed topological string partition functions. For example, our examples indicate that the deformed topological string partition functions satisfy a deformed Gopakumar-Vafa integrality. We will also show that many results in [23] and [12] can be generalized to the deformed case, i.e., we obtain some explicit expressions for some deformed topological string partition functions which we identify with the equivariant indices of some natural bundles on the framed moduli spaces.

The rest of the paper is arranged as follows. In Section 2 we compute the equivariant indices of some natural equivariant bundles on the framed moduli spaces. In Section 3 we collect some results on skew Schur functions which serve as our main technical tools. In Section 4 we will introduce the deformed topological vertex by studying the large N Chern-Simons invariants of the Hopf link. We present some product expressions for certain sums of these invariants in Section 5. We propose in Section 6 partition functions of some local Calabi-Yau spaces using the deformed topological vertex and identify with the equivariant indices in Section 2.

2. THE DEFORMED INSTANTON COUNTING

In this section we consider some equivariant indices of equivariant bundles extending the ones considered in [12]. They give us the “deformed instanton counting functions” to be identified with the “deformed curved counting functions” in §6.

2.1. The framed moduli spaces. Let $M(N, k)$ denote the framed moduli space of torsion free sheaves on $\mathbb{P}^2$ with rank N and $c_2 = k$. The framing means a trivialization of the sheaf restricted to the line at infinity. In particular when $N = 1$ the framed moduli spaces are the Hilbert schemes $\mathbb{C}^{[k]}$. See [16] for details.

As proved in [16], $M(N, k)$ is a nonsingular variety of dimension $2Nk$. The action of the maximal torus T of $GL_N(\mathbb{C})$ together with the torus action on $\mathbb{P}^2$ induces an action on $M(N, k)$. As shown in [16], the fixed points are isolated and parameterized by N -tuples of partitions $\vec{\mu} = (\mu^1, \dots, \mu^N)$ such that $\sum_i |\mu^i| = k$. The weight decomposition of the tangent bundle of $TM(N, k)$ at a fixed point $\vec{\mu}$ is given by

$$(1) \quad \sum_{\alpha, \gamma=1}^N e_\gamma e_\alpha^{-1} \left(\sum_{(i,j) \in \mu^\alpha} t_1^{-((\mu^\gamma)_j^t - i)} t_2^{\mu_i^\alpha - j + 1} + \sum_{(i,j) \in \mu^\gamma} t_1^{(\mu^\alpha)_j^t - i + 1} t_2^{-(\mu_i^\gamma - j)} \right),$$

where $t_1, t_2 \in \mathbb{C}^* \times \mathbb{C}^*$, and $e_\alpha \in T$.

The space $M(N, k)$ has the following remarkable property. Let E be an equivariant coherent sheaf on it. Even though $H^i(M(N, k), E)$ might be infinite-dimensional, it still makes sense to define the equivariant index and compute it by localization (cf. [16]):

Lemma 2.1. *Let E be an equivariant coherent sheaf on $M(N, k)$. Then*

$$\chi(M(N, k), E) = \sum_{i=0}^{2Nk} (-1)^i \operatorname{ch} H^i(M(N, k), E) = \sum_{\vec{\mu}} \operatorname{ch} \left(\frac{i_{\vec{\mu}}^* E}{\wedge_{-1} T_{\vec{\mu}}^* M(N, k)} \right).$$

2.2. Some naturally defined bundles on the framed moduli spaces. Recall $M(N, k)$ can be identified with the space of equivalent classes of tuples of linear maps

$$(B_1 : V \rightarrow V; B_2 : V \rightarrow V; i : W \rightarrow V; j : V \rightarrow W)$$

satisfying

$$[B_1, B_2] + ij = 0$$

and a stability condition. Hence one gets a vector bundle $\mathbb{V}_k$ over $M(N, k)$ whose fibers are given by V . This bundle is an equivariant bundle, and its weight decomposition at a fixed point $\vec{\mu}$ is given by [16]:

$$(2) \quad \mathbb{V}_k = \bigoplus_{\alpha} e_{\alpha} \sum_{(i,j) \in \mu^\alpha} t_1^{-i+1} t_2^{-j+1}.$$

Therefore, the weight of $\det \mathbb{V}_k^*$ at the fixed point $\vec{\mu}$ is

$$\prod_{\alpha} \left(e_{\alpha}^{-|\mu^\alpha|} \prod_{(i,j) \in \mu^\alpha} t_1^{i-1} t_2^{j-1} \right).$$

One can has an equivariant bundle $\mathbb{W}_k$ whose fibers are given by W . It has the following weight decomposition:

$$(3) \quad \mathbb{W}_k = \sum_{\alpha=1}^N e_{\alpha}.$$

Now we take $E_{N,k}^m = K_{N,k}^{\frac{1}{2}} \otimes (\det \mathbb{V}_k^*)^m$, where $K_{N,k}$ denotes the canonical line bundle of $M(N, k)$. By Lemma 2.1, (1), (2) and (3) one easily gets

Lemma 2.2. *We have*

$$(4) \quad \begin{aligned} & \sum_{k \geq 0} Q^k \chi(M(N, k), \Lambda_{-e^{-t}}(\mathbb{V}_k \otimes \mathbb{W}_k^*) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_k^* \otimes \mathbb{W}_k) \otimes E_{N,k}^m)(e_1, \dots, e_N, t_1, t_2) \\ &= \sum_{\mu^1, \dots, \mu^N} Q^{\sum_{i=1}^N |\mu^i|} \prod_{\alpha=1}^N \left(e_{\alpha}^{-|\mu^{\alpha}|} \prod_{(i,j) \in \mu^{\alpha}} t_1^{i-1} t_2^{j-1} \right)^m \\ & \cdot \prod_{\alpha, \gamma=1}^N \prod_{(i,j) \in \mu^{\alpha}} \frac{1 - e^{-t} e_{\alpha} e_{\gamma}^{-1} t_1^{-i+1} t_2^{-j+1}}{(e_{\alpha}^{-1} e_{\gamma} t_1^{-(\mu^{\gamma})_j^t - i} t_2^{\mu_i^{\alpha} - j + 1})^{\frac{1}{2}} - (e_{\alpha}^{-1} e_{\gamma} t_1^{-(\mu^{\gamma})_j^t - i} t_2^{\mu_i^{\alpha} - j + 1})^{-\frac{1}{2}}} \\ & \cdot \prod_{(i,j) \in \mu^{\gamma}} \frac{1 - e^{-t} (e_{\gamma} e_{\alpha}^{-1} t_1^{-i+1} t_2^{-j+1})^{-1}}{(e_{\alpha}^{-1} e_{\gamma} t_1^{(\mu^{\alpha})_j^t - i + 1} t_2^{-(\mu_i^{\gamma} - j)})^{\frac{1}{2}} - (e_{\alpha}^{-1} e_{\gamma} t_1^{(\mu^{\alpha})_j^t - i + 1} t_2^{-(\mu_i^{\gamma} - j)})^{-\frac{1}{2}}}. \end{aligned}$$

2.3. Rank 1 case and generalizations. When $N = 1$, $t_1 = q$ and $t_2 = q^{-1}$, we have

$$(5) \quad \begin{aligned} & \sum_{k \geq 0} Q^k \chi((\mathbb{C}^2)^{[k]}, \Lambda_{-e^{-t}}(\mathbb{V}_k \otimes \mathbb{W}_k^*) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_k^* \otimes \mathbb{W}_k) \otimes K_{1,k})(q, q^{-1}) \\ &= \sum_{\mu} Q^{|\mu|} \prod_{x \in \mu} \frac{1 - e^{-t} q^{c(x)}}{q^{-h(x)/2} - q^{h(x)/2}} \cdot \frac{1 - e^{-t} q^{-c(x)}}{q^{h(x)/2} - q^{-h(x)/2}}. \end{aligned}$$

Since the canonical line bundle $K_{1,k}$ is equivariantly trivial, this is equal to:

$$(6) \quad \begin{aligned} & \sum_{k \geq 0} Q^k \chi(M(1, k), \Lambda_{-e^{-t}}(\mathbb{V}_k \otimes \mathbb{W}_k^*) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_k^* \otimes \mathbb{W}_k))(q, q^{-1}) \\ &= \sum_{\mu} Q^{|\mu|} \prod_{x \in \mu} \left(\frac{1 - e^{-t} q^{c(x)}}{1 - q^{h(x)}} \cdot \frac{1 - e^{-t} q^{-c(x)}}{1 - q^{-h(x)}} \right). \end{aligned}$$

This can be generalized as follows. Let

$$W_{e^{-t}, y}(\mathbb{V}_n) = \Lambda_{-e^{-t}}(\mathbb{V}_n) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_n^*) \otimes S_{e^{-t}y}(\mathbb{V}_n) \otimes S_{e^{-t}y}(\mathbb{V}_n^*).$$

Now by (1) and (2),

$$\begin{aligned}
& \chi((\mathbb{C}^2)^{[n]}, \Lambda_{-y}(T^*(\mathbb{C}^2)^{[n]}) \otimes W_{e^{-t}, y}(\mathbb{V}_n))(t_1, t_2) \\
(7) \quad &= \sum_{|\mu|=n} \prod_{(i,j) \in \mu} \frac{(1 - e^{-t} t_1^{-i+1} t_2^{-j+1})(1 - e^{-t} t_1^{i-1} t_2^{j-1})}{(1 - t_1^{\mu_j^t - i} t_2^{-(\mu_i - j + 1)})(1 - t_1^{-(\mu_j^t - i + 1)} t_2^{\mu_i - j})} \\
&\quad \cdot \prod_{(i,j) \in \mu} \frac{(1 - y t_1^{\mu_j^t - i} t_2^{-(\mu_i - j + 1)})(1 - y t_1^{-(\mu_j^t - i + 1)} t_2^{\mu_i - j})}{(1 - y e^{-t} t_1^{-i+1} t_2^{-j+1})(1 - y e^{-t} t_1^{i-1} t_2^{j-1})}.
\end{aligned}$$

This generalizes the equivariant χ_y genera of the Hilbert schemes as one can see by taking $e^{-t} = 0$. It is easy to see that

$$\begin{aligned}
& \chi((\mathbb{C}^2)^{[n]}, \Lambda_{-y}(T^*(\mathbb{C}^2)^{[n]}) \otimes W_{e^{-t}, y}(\mathbb{V}_n))(q, q^{-1}) \\
(8) \quad &= \sum_{|\mu|=n} \prod_{x \in \mu} \left(\frac{(1 - e^{-t} q^{c(x)})(1 - e^{-t} q^{-c(x)})}{(1 - y e^{-t} q^{c(x)})(1 - y e^{-t} q^{-c(x)})} \frac{(1 - y q^{h(x)})(1 - y q^{-h(x)})}{(1 - q^{h(x)})(1 - q^{-h(x)})} \right).
\end{aligned}$$

We will generalize this further to an analogue of the elliptic genus as follows. Define

$$\begin{aligned}
\text{Ell}(T(\mathbb{C}^2)^{[n]}, \mathbb{V}_n)(y, p, e^{-t}) &= \bigotimes_{n=1}^{\infty} (\Lambda_{-y p^{n-1}}(T^*(\mathbb{C}^2)^{[n]}) \otimes \Lambda_{-y^{-1} p^n}(T(\mathbb{C}^2)^{[n]}) \\
&\quad \otimes S_{p^n}(T^*(\mathbb{C}^2)^{[n]}) \otimes S_{p^n}(T(\mathbb{C}^2)^{[n]}) \\
&\quad \otimes \Lambda_{-e^{-t} p^{n-1}}(\mathbb{V}_n \oplus \mathbb{V}_n^*) \otimes \Lambda_{-e^{-t} p^n}(\mathbb{V}_n \oplus \mathbb{V}_n^*)^* \\
&\quad \otimes S_{e^{-t} y p^{n-1}}(\mathbb{V}_n \oplus \mathbb{V}_n^*) \otimes S_{e^{-t} y^{-1} p^n}(\mathbb{V}_n \oplus \mathbb{V}_n^*)^*.
\end{aligned}$$

This suggests that we should consider the supermanifolds obtained from $(\mathbb{C}^2)^{[n]}$ whose odd part of the tangent spaces are $\mathbb{V}_n \oplus \mathbb{V}_n^*$. By taking $p = 0$, one gets:

$$\begin{aligned}
& \text{Ell}(T(\mathbb{C}^2)^{[n]}, \mathbb{V}_n)(y, 0, e^{-t}) \\
&= \Lambda_{-y}(T^*(\mathbb{C}^2)^{[n]}) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_n) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_n^*) \otimes S_{e^{-t} y}(\mathbb{V}_n) \otimes S_{e^{-t} y}(\mathbb{V}_n^*) \\
&= \Lambda_{-y}(T^*(\mathbb{C}^2)^{[n]}) \otimes W_{e^{-t}, y}(\mathbb{V}_n).
\end{aligned}$$

by (1) and (2),

$$\begin{aligned}
& \chi((\mathbb{C}^2)^{[n]}, \text{Ell}(T(\mathbb{C}^2)^{[n]}, \mathbb{V}_n)(y, p, e^{-t}))(t_1, t_2) \\
(9) \quad &= \sum_{|\mu|=n} \prod_{(i,j) \in \mu} \prod_{k=1}^{\infty} \left(\frac{(1 - y p^{k-1} t_1^{\mu_j^t - i} t_2^{-(\mu_i - j + 1)})(1 - y p^{k-1} t_1^{-(\mu_j^t - i + 1)} t_2^{\mu_i - j})}{(1 - p^{k-1} t_1^{\mu_j^t - i} t_2^{-(\mu_i - j + 1)})(1 - p^{k-1} t_1^{-(\mu_j^t - i + 1)} t_2^{\mu_i - j})} \right. \\
&\quad \cdot \frac{(1 - y p^k t_1^{-(\mu_j^t - i)} t_2^{\mu_i - j + 1})(1 - y p^k t_1^{\mu_j^t - i + 1} t_2^{-(\mu_i - j)})}{(1 - p^k t_1^{-(\mu_j^t - i)} t_2^{\mu_i - j + 1})(1 - p^k t_1^{\mu_j^t - i + 1} t_2^{-(\mu_i - j)})} \\
&\quad \cdot \frac{(1 - e^{-t} p^{k-1} t_1^{-i+1} t_2^{-j+1})(1 - e^{-t} p^{k-1} t_1^{i-1} t_2^{j-1})}{(1 - y e^{-t} p^{k-1} t_1^{-i+1} t_2^{-j+1})(1 - y e^{-t} p^{k-1} t_1^{i-1} t_2^{j-1})} \\
&\quad \left. \cdot \frac{(1 - e^{-t} p^k t_1^{-i+1} t_2^{-j+1})(1 - e^{-t} p^k t_1^{i-1} t_2^{j-1})}{(1 - y e^{-t} p^k t_1^{-i+1} t_2^{-j+1})(1 - y e^{-t} p^k t_1^{i-1} t_2^{j-1})} \right).
\end{aligned}$$

It follows easily that

$$\begin{aligned}
 & \chi((\mathbb{C}^2)^{[n]}, \text{Ell}(T(\mathbb{C}^2)^{[n]}, \mathbb{V}_n)(y, p, e^{-t}))(q, q^{-1}) \\
 &= \sum_{|\mu|=n} \prod_{(i,j) \in \mu} \prod_{k=1}^{\infty} \left(\frac{(1 - p^{k-1} y q^{h(i,j)})(1 - p^{k-1} y q^{-h(i,j)})}{(1 - p^{k-1} q^{h(i,j)})(1 - p^{k-1} q^{-h(i,j)})} \right. \\
 (10) \quad & \cdot \frac{(1 - p^k y^{-1} q^{h(i,j)})(1 - p^k y^{-1} q^{-h(i,j)})}{(1 - p^k q^{h(i,j)})(1 - p^k q^{-h(i,j)})} \\
 & \cdot \frac{(1 - e^{-t} p^{k-1} q^{c(i,j)})(1 - e^{-t} p^{k-1} q^{-c(i,j)})}{(1 - e^{-t} p^{k-1} y q^{c(i,j)})(1 - e^{-t} p^{k-1} y q^{-c(i,j)})} \\
 & \cdot \left. \frac{(1 - e^{-t} p^k q^{c(i,j)})(1 - e^{-t} p^k q^{-c(i,j)})}{(1 - e^{-t} p^k y^{-1} q^{c(i,j)})(1 - e^{-t} p^k y^{-1} q^{-c(i,j)})} \right).
 \end{aligned}$$

It is natural to expect that a deformed version of the equivariant DMVV conjecture holds for such deformed elliptic genera.

2.4. Rank > 1 case. When $t_1 = q$, $t_2 = q^{-1}$, after a tedious elementary calculation one can find:

$$\begin{aligned}
 (11) \quad & \sum_{k \geq 0} Q^k \chi(M(N, k), \Lambda_{-e^{-t}}(\mathbb{V}_k \otimes \mathbb{W}_k^*) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_k^* \otimes \mathbb{W}_k) \otimes E_{N,k}^m)(e_1, \dots, e_N, q, q^{-1}) \\
 &= \sum_{\mu^1, \dots, \mu^N} Q^{\sum_{i=1}^N |\mu^i|} e^{-Nt \sum_{\alpha=1}^N |\mu^\alpha|} \prod_{\alpha=1}^N \left(e_\alpha^{-|\mu^\alpha|} q^{-\kappa_{\mu^\alpha}/2} \right)^m \\
 & \cdot \prod_{\alpha=1}^N \prod_{(i,j) \in \mu^\alpha} \frac{e^{t/2} q^{(j-i)/2} - e^{-t/2} q^{-(j-i)/2}}{q^{h(i,j)/2} - q^{-h(i,j)/2}} \\
 & \cdot \prod_{1 \leq \alpha < \gamma \leq N} \left(\prod_{(i,j) \in \mu^\alpha} \frac{1 - e^{-t} e_\alpha e_\gamma^{-1} q^{j-i}}{1 - e_\alpha e_\gamma^{-1} q^{\mu_i^\alpha + (\mu^\gamma)_j^t - i - j + 1}} \right. \\
 & \cdot \prod_{(i,j) \in \mu^\gamma} \frac{1 - e^{-t} e_\alpha e_\gamma^{-1} q^{-(j-i)}}{1 - e_\alpha e_\gamma^{-1} q^{-(\mu_i^\gamma + (\mu^\alpha)_j^t - i - j + 1)}} \Big) \\
 & \times \prod_{\alpha=1}^N \prod_{(i,j) \in \mu^\alpha} \frac{e^{-t/2} q^{(j-i)/2} - e^{t/2} q^{-(j-i)/2}}{q^{h(i,j)/2} - q^{-h(i,j)/2}} \\
 & \cdot \prod_{1 \leq \alpha < \gamma \leq N} \left(\prod_{(i,j) \in \mu^\alpha} \frac{1 - e^t e_\alpha e_\gamma^{-1} q^{j-i}}{1 - e_\alpha e_\gamma^{-1} q^{\mu_i^\alpha + (\mu^\gamma)_j^t - i - j + 1}} \right. \\
 & \cdot \prod_{(i,j) \in \mu^\gamma} \frac{1 - e^t e_\alpha e_\gamma^{-1} q^{-(j-i)}}{1 - e_\alpha e_\gamma^{-1} q^{-(\mu_i^\gamma + (\mu^\alpha)_j^t - i - j + 1)}} \Big).
 \end{aligned}$$

3. PRELIMINARY RESULTS ON SKEW SCHUR FUNCTIONS

In this section we collect some results on skew Schur functions. They will be our main technical tools for the rest of the paper.

3.1. Summation formulas for Skew Schur functions. Recall the following identity:

$$(12) \quad \prod_{i,j=1}^{\infty} (1 - Qx_i y_j)^{-1} = \exp \sum_{n=1}^{\infty} \frac{Q^n}{n} p_n(x) p_n(y),$$

where $x = (x_1, x_2, \dots)$, $y = (y_1, y_2, \dots)$, and

$$p_n(x) = x_1^n + x_2^n + \dots.$$

For some specializations, the left-hand side of (12) may not make sense, but the right-hand side makes sense. For example, when $x = q^\rho = (q^{-\frac{1}{2}}, q^{-\frac{3}{2}}, \dots)$, $y = q^{-\rho} = (q^{\frac{1}{2}}, q^{\frac{3}{2}}, \dots)$, the left-hand side becomes

$$\prod_{i,j=1}^{\infty} (1 - Qq^{j-i}),$$

which does not make sense, while on the right-hand,

$$(13) \quad p_n(q^\rho) = q^{-\frac{n}{2}} + q^{-\frac{3n}{2}} + \dots = \frac{q^{-\frac{n}{2}}}{1 - q^{-n}} = -\frac{q^{\frac{n}{2}}}{1 - q^n} = -p_n(q^{-\rho}),$$

hence the right-hand side of (12) is

$$\exp \left(- \sum_{n=1}^{\infty} \frac{(qQ)^n}{(q^n - 1)^2} \right).$$

We will use the following identity for skew Schur functions (cf. [13, p. 93, (1)]):

$$(14) \quad \sum_{\eta} s_{\eta/\mu}(x) s_{\eta/\nu}(y) = \exp \sum_{n=1}^{\infty} \frac{1}{n} p_n(x) p_n(y) \cdot \sum_{\tau} s_{\mu/\tau}(y) s_{\nu/\tau}(x),$$

$$(15) \quad \sum_{\eta} s_{\eta/\mu^t}(x) s_{\eta^t/\nu}(y) = \exp \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} p_n(x) p_n(y) \cdot \sum_{\tau} s_{\mu/\tau^t}(y) s_{\nu^t/\tau}(x),$$

where $x = (x_1, x_2, \dots)$, $y = (y_1, y_2, \dots)$. In particular,

$$(16) \quad \sum_{\eta} s_{\eta}(x) s_{\eta}(y) = \exp \sum_{n=1}^{\infty} \frac{1}{n} p_n(x) p_n(y),$$

$$(17) \quad \sum_{\eta} s_{\eta/\mu}(x) s_{\eta}(y) = \exp \sum_{n=1}^{\infty} \frac{1}{n} p_n(x) p_n(y) \cdot s_{\mu}(y),$$

$$(18) \quad \sum_{\eta} s_{\eta}(x) s_{\eta^t}(y) = \exp \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} p_n(x) p_n(y),$$

$$(19) \quad \sum_{\eta} s_{\eta/\mu^t}(x) s_{\eta}(y) = \exp \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} p_n(x) p_n(y) \cdot s_{\mu}(y).$$

For $x = (x_1, x_2, \dots)$, $y = (y_1, y_2, \dots)$, let $(x, y) = (x_1, y_1, x_2, y_2, \dots)$. Then we have

$$(20) \quad \sum_{\xi} s_{\mu/\xi}(x) s_{\xi/\nu}(y) = s_{\mu/\nu}(x, y).$$

We will need the following:

Lemma 3.1. [23] *The following identity holds:*

$$(21) \quad \sum_{\nu^1, \dots, \nu^N} \sum_{\eta^1, \dots, \eta^{N-1}} \prod_{k=1}^N s_{\nu^k / \eta^{k-1}}(x^k) Q_k^{|\nu^k|} s_{\nu^k / \eta^k}(y^k) \\ = \prod_{1 \leq k < l \leq N+1} \exp \sum_{n=1}^{\infty} \frac{(Q_k Q_{k+1} \cdots Q_{l-1})^n}{n} p_n(x^k) p_n(y^{l-1}),$$

where $\eta^0 = \eta^N = (0)$, $x = (x_1^k, x_2^k, \dots)$, $y^k = (y_1^k, y_2^k, \dots)$.

Formulas of the type in the following Lemma appeared in [7] without proof.

Lemma 3.2. *For $|uv| < 1$ we have*

$$(22) \quad \sum_{\mu, \nu} u^{|\mu|} v^{|\nu|} s_{\mu/\nu}(x) s_{\mu/\nu}(y) = \exp \sum_{n=1}^{\infty} \frac{u^n}{n(1 - (uv)^n)} p_n(x) p_n(y),$$

$$(23) \quad \sum_{\mu, \nu} (-u)^{|\mu|} (-v)^{|\nu|} s_{\mu/\nu^t}(x) s_{\mu^t/\nu}(y) = \exp \sum_{n=1}^{\infty} \frac{-u^n}{n(1 - (uv)^n)} p_n(x) p_n(y).$$

For $|u_1 u_2 v_1 v_2| < 1$, we have

$$(24) \quad \sum_{\mu, \nu, \xi, \eta} u_1^{|\mu|} u_2^{|\nu|} v_1^{|\xi|} v_2^{|\eta|} s_{\mu/\xi}(x) s_{\nu/\xi}(y) s_{\nu/\eta}(z) s_{\mu/\eta}(w) \\ = \exp \sum_{n=1}^{\infty} \frac{1}{n(1 - (u_1 u_2 v_1 v_2)^n)} (u_1^n p_n(x) p_n(z) + u_2^n p_n(y) p_n(w) \\ + (u_1 u_2 v_1)^n p_n(w) p_n(z) + (u_1 u_2 v_2)^n p_n(x) p_n(y)),$$


and

$$(25) \quad \sum_{\mu, \nu, \xi, \eta} (-u_1)^{|\mu|} (-u_2)^{|\nu|} v_1^{|\xi|} v_2^{|\eta|} s_{\mu/\xi^t}(x) s_{\nu/\xi^t}(y) s_{\nu^t/\eta}(z) s_{\mu^t/\eta}(w) \\ = \exp \sum_{n=1}^{\infty} \frac{1}{n(1 - (u_1 u_2 v_1 v_2)^n)} (-u_1^n p_n(x) p_n(z) - u_2^n p_n(y) p_n(w) \\ + (u_1 u_2 v_1)^n p_n(w) p_n(z) + (u_1 u_2 v_2)^n p_n(x) p_n(y)),$$

Proof. Write the left-hand side of (22) as $S(x, y, u, v)$. By (15) one has:

$$\begin{aligned} S(x, y; u, v) &= \sum_{\nu} (uv)^{|\nu|} \sum_{\mu} s_{\mu/\nu}(ux) s_{\mu/\nu}(y) \\ &= \exp \sum_{n=1}^{\infty} \frac{u^n}{n} p_n(x) p_n(y) \cdot \sum_{\nu, \mu} (uv)^{|\nu|} s_{\nu/\mu}(ux) s_{\nu/\mu}(y) \\ &= \exp \sum_{n=1}^{\infty} \frac{u^n}{n} p_n(x) p_n(y) \cdot \sum_{\nu, \mu} v^{|\nu|} u^{|\mu|} s_{\nu/\mu}(vy) s_{\nu/\mu}(ux) \\ &= \exp \sum_{n=1}^{\infty} \frac{u^n}{n} p_n(x) p_n(y) \cdot S(vy, ux; u, v). \end{aligned}$$

This procedure can be graphically represented as follows:



One can repeat this procedure for infinitely many times to get:

$$\begin{aligned}
& \sum_{\mu, \nu} u^{|\mu|} v^{|\nu|} s_{\mu/\nu}(x) s_{\mu/\nu}(y) \\
&= \exp \sum_{n=1}^{\infty} \frac{1 + u^n + u^{2n} + (uv)^{2mn}}{n} p_n(x) p_n(y) \cdot S((uv)^m x, (uv)^m y; u) \\
&= \exp \sum_{n=1}^{\infty} \frac{u^n (1 + (uv)^n + \dots + (uv)^{2n} + \dots)}{n} p_n(x) p_n(y) \\
&= \exp \sum_{n=1}^{\infty} \frac{u^n}{n(1 - (uv)^n)} p_n(x) p_n(y).
\end{aligned}$$

The proof of (23) is similar. We present two proof for (24). In the first proof we use (14) and (20) to reduce to (22):

$$\begin{aligned}
& \sum_{\mu, \nu, \xi, \eta} u_1^{|\mu|} u_2^{|\nu|} v_1^{|\xi|} v_2^{|\eta|} s_{\mu/\xi}(x) s_{\nu/\xi}(y) s_{\mu/\eta}(z) s_{\nu/\eta}(w) \\
&= \sum_{\mu, \xi, \eta} u_1^{|\mu|} u_2^{|\xi|} v_1^{|\xi|} v_2^{|\eta|} s_{\mu/\xi}(x) s_{\mu/\eta}(z) \sum_{\nu} s_{\nu/\xi}(u_2 y) s_{\nu/\eta}(w) \\
&= \exp \sum_{n=1}^{\infty} \frac{u_2^n}{n} p_n(y) p_n(w) \cdot \sum_{\mu, \nu, \xi, \eta} u_1^{|\mu|} u_2^{|\xi|} v_1^{|\xi|} v_2^{|\eta|} s_{\mu/\xi}(x) s_{\mu/\eta}(z) s_{\eta/\nu}(u_2 y) s_{\xi/\nu}(w) \\
&= \exp \sum_{n=1}^{\infty} \frac{u_2^n}{n} p_n(y) p_n(w) \cdot \sum_{\mu, \nu} u_1^{|\mu|} (u_2 v_1 v_2)^{|\nu|} s_{\mu/\nu}(x, u_2 v_1 w) s_{\mu/\nu}(z, u_2 v_2 y) \\
&= \exp \sum_{n=1}^{\infty} \frac{u_2^n}{n} p_n(y) p_n(w) \cdot \exp \sum_{n=1}^{\infty} \frac{u_1^n}{n(1 - (u_1 u_2 v_1 v_2)^n)} (p_n(x) \\
&\quad + (u_2 v_1)^n p_n(w)) (p_n(z) + (u_2 v_2)^n p_n(y)) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n(1 - (u_1 u_2 v_1 v_2)^n)} (u_1^n p_n(x) p_n(z) \\
&\quad + (u_1 u_2 v_2)^n p_n(x) p_n(y) + (u_1 u_2 v_1)^n p_n(w) p_n(z) + u_2^n p_n(y) p_n(w))
\end{aligned}$$

The second proof of (24) is similar to the proof of (22). We write its left-hand side as $S(x, y, z, w; u, v)$. Then we have:

$$\begin{aligned}
& S(x, y, z, w; u_1, u_2, v_1, v_2) \\
&= \sum_{\mu, \nu, \xi, \eta} u_1^{|\mu|} u_2^{|\nu|} v_1^{|\xi|} v_2^{|\eta|} s_{\mu/\xi}(x) s_{\nu/\eta}(y) s_{\mu/\eta}(z) \\
&= \sum_{\xi, \eta} (u_1 v_1)^{|\xi|} (u_2 v_2)^{|\eta|} \sum_{\mu} s_{\mu/\xi}(u_1 x) s_{\mu/\eta}(z) \sum_{\nu} s_{\nu/\xi}(y) s_{\nu/\eta}(u_2 w) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n} (u_1^n p_n(x) p_n(z) + u_2^n p_n(y) p_n(w)) \\
&\quad \cdot \sum_{\mu, \nu, \xi, \eta} (u_1 v_1)^{|\xi|} (u_2 v_2)^{|\eta|} s_{\eta/\mu}(u_1 x) s_{\xi/\mu}(z) s_{\eta/\nu}(y) s_{\xi/\nu}(u_2 w) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n} (u_1^n p_n(x) p_n(z) + u_2^n p_n(y) p_n(w)) \\
&\quad \cdot \sum_{\mu, \nu, \xi, \eta} u_1^{|\xi|} v_1^{|\mu|} u_2^{|\eta|} v_2^{|\nu|} s_{\eta/\mu}(u_1 x) s_{\xi/\mu}(v_1 z) s_{\eta/\nu}(v_2 y) s_{\xi/\nu}(u_2 w) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n} (u_1^n p_n(x) p_n(z) + u_2^n p_n(y) p_n(w)) \cdot S(v_1 z, u_1 x, u_2 w, v_2 y; u_1, u_2, v_1, v_2).
\end{aligned}$$

Repeating this procedure for four times:

$$\begin{aligned}
& S(x, y, z, w; u_1, u_2, v_1, v_2) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n} (u_1^n (1 + (u_1 u_2 v_1 v_2)^n) p_n(x) p_n(z) + u_2^n (1 + (u_1 u_2 v_1 v_2)^n) p_n(y) p_n(w) \\
&\quad + (u_1 u_2 v_1)^n (1 + (u_1 u_2 v_1 v_2)^n) p_n(w) p_n(z) \\
&\quad + (u_1 u_2 v_2)^n (1 + (u_1 u_2 v_1 v_2)^n) p_n(x) p_n(y)) \\
&\quad \cdot S(u_1 u_2 v_1 v_2 x, u_1 u_2 v_1 v_2 y, u_1 u_2 v_1 v_2 z, u_1 u_2 v_1 v_2 w; u_1, u_2, v_1, v_2).
\end{aligned}$$

Repeating this procedure for infinitely many times:

$$\begin{aligned}
& S(x, y, z, w; u_1, u_2, v_1, v_2) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n(1 - (u_1 u_2 v_1 v_2)^n)} (u_1^n p_n(x) p_n(z) + u_2^n p_n(y) p_n(w) \\
&\quad + (u_1 u_2 v_1)^n p_n(w) p_n(z) + (u_1 u_2 v_2)^n p_n(x) p_n(y)).
\end{aligned}$$

□

3.2. Some results on specialization of Schur functions. Recall any symmetric function f can be written a polynomial $f(e_1, e_2, \dots, e_n, \dots)$ in the elementary symmetric functions $e_1, \dots, e_n, \dots$. Let $E(u) = 1 + \sum_{n=1}^{\infty} e_n u^n$. We write $f(e_1, \dots, e_n, \dots)$ as $f(E(u))$. In general, take $E(u)$ to be any formal power series with leading coefficient 1, $f(E(u))$ defines a specialization of f .

Lemma 3.3. *Let $a = (a_1, a_2, \dots)$, $b = (b_1, b_2, \dots)$, and*

$$E(u) = \frac{\prod_{i=1}^{\infty} (1 + a_i u)}{\prod_{i=1}^{\infty} (1 + b_i u)}.$$

Then we have

$$(26) \quad p_n(E(u)) = p_n(a) - p_n(b)$$

Proof. Let

$$P(u) = \sum_{n=1}^{\infty} p_n(E(u)) u^{n-1}.$$

Then one has [13]:

$$\begin{aligned} P(-u) &= \frac{d}{du} \log E(u) = \sum_{i=1}^{\infty} \left(\frac{a_i}{1+a_i u} - \frac{b_i}{1+b_i u} \right) \\ &= \sum_{i=1}^{\infty} \sum_{n=1}^{\infty} (a_i^n - b_i^n) (-u)^{n-1} = \sum_{n=1}^{\infty} (p_n(a) - p_n(b)) (-u)^{n-1}. \end{aligned}$$

□

Lemma 3.4. Let $a = (a_1, a_2, \dots)$, $b = (b_1, b_2, \dots)$, $c = (c_1, c_2, \dots)$, $d = (d_1, \dots)$, and

$$E(u) = \frac{\prod_{i=1}^{\infty} (1 + a_i u)}{\prod_{i=1}^{\infty} (1 + b_i u)}, \quad \tilde{E}(u) = \frac{\prod_{i=1}^{\infty} (1 + c_i u)}{\prod_{i=1}^{\infty} (1 + d_i u)}.$$

Then we have

$$(27) \quad \sum_{\mu} s_{\mu}(E(u)) s_{\mu}(\tilde{E}(u)) = \prod_{i,j=1}^{\infty} \frac{(1 - a_i d_j)(1 - b_i c_j)}{(1 - a_i c_j)(1 - b_i d_j)}.$$

Proof. This is a straightforward consequence of (16) and Lemma 3.3. □

4. THE DEFORMED TOPOLOGICAL VERTEX

In this section we will introduce the deformed topological vertex. We will begin by studying the large N Chern-Simons invariants of the Hopf link. We will express them in terms of specializations of skew Schur functions and study their symmetric properties. It will be interesting to combine our approach with that of Awata and Konno [4]. We hope to report on this in a separate work.

4.1. The quantum dimension. Recall the large N invariant of the unknot is given by the quantum dimension [14]:

$$(28) \quad W_{\mu}(q, e^{-t}) = \prod_{e \in \mu} \frac{e^{t/2} q^{c(e)/2} - e^{-t/2} q^{-c(e)/2}}{q^{h(e)/2} - q^{-h(e)/2}},$$

where $\mu = (\mu_1, \dots, \mu_l)$ is a partition. For the zero partition we take $W_{(0)}(q, e^{-t}) = 1$. Here $c(e)$ and $h(e)$ are the content and the hook length of the box e in the Young diagram μ respectively. From the definition (28) one easily gets the following:

Proposition 4.1. *The quantum dimension has the following symmetry properties:*

$$(29) \quad W_{\mu}(q^{-1}, e^t) = W_{\mu}(q, e^{-t}),$$

$$(30) \quad W_{\mu^t}(q^{-1}, e^{-t}) = W_{\mu^t}(q, e^t) = (-1)^{|\mu|} W_{\mu}(q, e^{-t}).$$

Our next result express the quantum dimension as a specialization of the Schur functions.

Proposition 4.2. *The quantum dimension can be identified with the following specialization of the Schur function:*

$$(31) \quad W_\mu(q, e^{-t}) = (qe^t)^{|\mu|/2} s_\mu(E^{(0)}(u; q, e^{-t})),$$

where

$$(32) \quad E^{(0)}(u) = 1 + \sum_{n=1}^{\infty} u^n e_n = 1 + \sum_{n=1}^{\infty} u^n \prod_{i=1}^n \frac{1 - e^{-t} q^{i-1}}{q^i - 1}.$$

Proof. Recall the following fact from the theory of symmetric functions. For a specialization with

$$H(u) = \sum_{r=0}^{\infty} h_r u^r = \prod_{i=0}^{\infty} \frac{1 - bq^i t}{1 - aq^i t},$$

one has ([13], p. 27, Example 5, and p. 45, Example 3).

$$(33) \quad h_r = \prod_{i=1}^r \frac{a - bq^{i-1}}{1 - q^i},$$

$$(34) \quad e_r = \prod_{i=1}^r \frac{aq^{i-1} - b}{1 - q^i},$$

$$(35) \quad p_r = \frac{a^r - b^r}{1 - q^r},$$

$$(36) \quad s_\nu = q^{n(\nu)} \prod_{x \in \nu} \frac{a - bq^{c(x)}}{1 - q^{h(x)}}.$$

Hence one gets $E(u)$ in (32) by taking $a = e^{-t}$, $b = 1$, and $q = e^{\sqrt{-1}\lambda}$. It follows that

$$\begin{aligned} s_\mu(E(u; q, e^{-t})) &= q^{n(\mu)} \prod_{e \in \mu} \frac{e^{-t} - q^{c(e)}}{1 - q^{h(e)}} \\ &= q^{n(\mu) + \frac{1}{2} \sum_{e \in \mu} [c(e) - h(e)]} e^{-|\mu|t/2} \prod_{e \in \mu} \frac{e^{t/2} q^{c(e)/2} - e^{-t/2} q^{-c(e)/2}}{q^{h(e)/2} - q^{-h(e)/2}} \\ &= (q^{-1} e^{-t})^{|\mu|/2} \prod_{e \in \mu} \frac{e^{t/2} q^{c(e)/2} - e^{-t/2} q^{-c(e)/2}}{q^{h(e)/2} - q^{-h(e)/2}} \\ &= (q^{-1} e^{-t})^{|\mu|/2} W_\mu(q, e^{-t}). \end{aligned}$$

Here we have used the following properties of the hook length and the content [13]:

$$(37) \quad \sum_{x \in \nu} h(x) = n(\nu) + n(\nu^t) + |\nu|,$$

$$(38) \quad \sum_{x \in \nu} c(x) = n(\nu^t) - n(\nu).$$

□

4.2. The invariant of the Hopf link. Recall the large N invariant of the Hopf link is [15, 2]:

$$(39) \quad W_{\mu,\nu}(q, e^{-t}) = W_{\mu}(q, e^{-t})(e^t q)^{|\nu|/2} s_{\nu}(E^{\mu}(u; q, e^{-t})),$$

where

$$E^{\mu}(u; q, e^{-t}) = \prod_{i=1}^{l(\mu)} \frac{1 + q^{\mu_i - i} u}{1 + q^{-i} u} \cdot (1 + \sum_{n=1}^{\infty} u^n \prod_{i=1}^n \frac{1 - e^{-t} q^{i-1}}{q^i - 1}).$$

Proposition 4.3. *We have*

$$(40) \quad E^{\mu}(u) = \frac{\prod_{i=1}^{\infty} (1 + q^{\mu_i - i} u)}{\prod_{i=1}^{\infty} (1 + e^{-t} q^{-i} u)}.$$

Proof. Recall the following famous identity due to Cauchy [3]. For $|x| < 1$, $|q| < 1$,

$$1 + \sum_{n=1}^{\infty} x^n \prod_{i=1}^n \frac{(1 - a q^{i-1})}{(1 - q^i)} = \prod_{i=0}^{\infty} \frac{(1 - a x q^i)}{(1 - x q^i)}.$$

Now

$$\begin{aligned} 1 + \sum_{n=1}^{\infty} u^n \prod_{i=1}^n \frac{1 - e^{-t} q^{i-1}}{q^i - 1} &= 1 + \sum_{n=1}^{\infty} (-e^{-t} q^{-1} u)^n \prod_{i=1}^n \frac{1 - e^t q^{-(i-1)}}{1 - q^{-i}} \\ &= \prod_{i=1}^{\infty} \frac{1 + q^{-i} u}{1 + e^{-t} q^{-i} u}. \end{aligned}$$

Hence (40) follows. $\square$

Apply Lemma 3.3 to $E^{\mu}(u)$:

$$(41) \quad p_n(E^{\mu}(u; q, e^{-t})) = \sum_{i=1}^{\infty} (q^{n(\mu_i - i)} - e^{-nt} q^{n(-i)}).$$

Now note

$$\sum_{i=1}^{\infty} q^{n(-i)} = \frac{q^{-n}}{1 - q^{-n}} = -\frac{1}{1 - q^n}.$$

Hence we have

$$(42) \quad p_n(E^{\mu}(u; q, e^{-t})) = p_n(q^{\mu_i + \rho - \frac{1}{2}}) + p_n(e^{-t} q^{-\rho - \frac{1}{2}}).$$

I.e., formally $E^{\mu}(u; q, e^{-t})$ gives the specialization $(q^{\mu + \rho - \frac{1}{2}}, e^{-t} q^{-\rho - \frac{1}{2}})$. In (42) and the expression below, we are not working in different domains $|q| < 1$ and $|q| > 1$ at the same time. Instead, we write the relevant skew Schur functions as polynomials of $p_n(q^{\rho})$ and $p_n(q^{-\rho})$, whose values are given by (13). Hence our treatment is mathematically rigorous. For later use note

$$(43) \quad p_n(E^{\mu}(u; q, e^{-t})) = p_1(E^{\mu}(q^n, e^{-nt})),$$

and

$$(44) \quad p_1(E^{\mu}(u; q, e^{-t})) = \sum_{i=1}^{\infty} (q^{\mu_i - i} - e^{-t} q^{-i}) = \sum_{i=1}^{l(\mu)} (q^{\mu_i - i} - q^{-i}) + \frac{1 - e^{-t}}{q - 1}.$$

Lemma 4.1. *We have*

$$(45) \quad s_\nu(E^\mu(u; q, e^{-t})) = q^{-|\nu|/2} \sum_{\eta} s_{\nu/\eta}(e^{-t} q^{-\rho}) s_\eta(q^{\mu+\rho})$$

$$(46) \quad = q^{-|\nu|/2} \sum_{\xi, \eta} s_{\nu/\eta}(e^{-t} q^{-\rho}) (-1)^{|\eta|} q^{\kappa_\eta/2} \frac{s_{\mu/\xi}(q^{-\rho}) s_{\eta/\xi}(q^{-\rho})}{s_\mu(q^{-\rho})}.$$

Proof. We use (42) and (20) to get:

$$\begin{aligned} s_\nu(E^\mu(u; q, e^{-t})) &= s_\nu(q^{\mu+\rho-\frac{1}{2}}, e^{-t} q^{-\rho-\frac{1}{2}}) = \sum_{\eta} s_{\nu/\eta}(e^{-t} q^{-\rho-\frac{1}{2}}) s_\eta(q^{\mu+\rho-\frac{1}{2}}) \\ &= \sum_{\eta} q^{-(|\nu|-|\eta|)/2} s_{\nu/\eta}(e^{-t} q^{-\rho}) q^{-|\eta|/2} s_\eta(q^{\mu+\rho}) \\ &= q^{-|\nu|/2} \sum_{\eta} s_{\nu/\eta}(e^{-t} q^{-\rho}) s_\eta(q^{\mu+\rho}). \end{aligned}$$

This proves (45). To prove (46) recall the following identity proved in [21]:

$$(47) \quad s_\nu(q^{\mu+\rho}) = (-1)^{|\mu|} q^{\kappa_\nu/2} \sum_{\xi} \frac{s_{\mu/\xi}(q^{-\rho}) s_{\nu/\xi}(q^{-\rho})}{s_\mu(q^{-\rho})}.$$

□

Corollary 4.1. *We have the following symmetry:*

$$(48) \quad s_{\nu^t/\eta^t}(E^{\mu^t}(u; q, e^{-t})) = (-1)^{|\nu|-|\eta|} q^{-(|\nu|-|\eta|)} s_{\nu/\eta}(E^\mu(u; q^{-1}, e^{-t})).$$

Proof. Recall the following symmetry [23]:

$$(49) \quad s_{\lambda/\mu}(q^{\nu+\rho}) = (-1)^{|\lambda|-|\mu|} s_{\lambda^t/\mu^t}(q^{-\nu-\rho}).$$

Hence (48) follows from (46). □

4.3. Symmetries of W_{μ^1, μ^2} .

Proposition 4.4. *We have*

$$(50) \quad W_{\mu^1, \mu^2}(q, e^{-t}) = W_{\mu^2, \mu^1}(q, e^{-t}),$$

$$(51) \quad W_{(\mu^1)^t, (\mu^2)^t}(q, e^{-t}) = (-1)^{|\mu^1|+|\mu^2|} W_{\mu^1, \mu^2}(q^{-1}, e^{-t}).$$

Proof. The identity (50) follows from the fact that $W_{\mu^1, \mu^2}(q, e^{-t})$ can be obtained from the colored HOMFLY polynomials of the Hopf link. By (39) and (??),

$$\begin{aligned} W_{\mu^t, \nu^t}(q, e^{-t}) &= W_{\mu^t}(q, e^{-t}) (e^t q)^{|\nu^t|/2} s_{\nu^t}(E_{\mu^t}(u; q, e^{-t})) \\ &= (-1)^{|\mu|} W_{\mu}(q^{-1}, e^{-t}) (e^t q)^{|\nu|/2} (-1)^{|\nu|} q^{-|\nu|} s_{\nu}(E_{\mu}(u; q^{-1}, e^{-t})) \\ &= (-1)^{|\mu|+|\nu|} W_{\mu, \nu}(q^{-1}, e^{-t}). \end{aligned}$$

This proves (51). □

4.4. The Deformed topological vertex. The topological vertex introduced in [1] is defined by

$$(52) \quad \mathcal{W}_{\mu^1, \mu^2, \mu^3}(q) = \sum_{\rho^1, \rho^3} c_{\rho^1(\rho^3)^t}^{\mu^1(\mu^3)^t} q^{\kappa_{\mu^2/2} + \kappa_{\mu^3/2}} \frac{\mathcal{W}_{(\mu^2)^t \rho^1}(q) \mathcal{W}_{\mu^2(\rho^3)^t}(q)}{\mathcal{W}_{\mu^2}(q)},$$

where

$$c_{\rho^1(\rho^3)^t}^{\mu^1(\mu^3)^t} = \sum_{\eta} c_{\eta \rho^1}^{\mu^1} c_{\eta(\rho^3)^t}^{(\mu^3)^t}.$$

It can be reformulated in terms of skew Schur functions [20, 23, 7]:

$$\mathcal{W}_{\mu^1, \mu^2, \mu^3}(q) = q^{\kappa_{\mu^3/2}} s_{\mu^2}(q^\rho) \sum_{\eta} s_{\mu^1/\eta}(q^{(\mu^2)^t + \rho}) s_{(\mu^3)^t/\eta}(q^{\mu^2 + \rho}).$$

Now we define the deformed topological vertex to be:

$$(53) \quad \begin{aligned} & W_{\mu^1, \mu^2, \mu^3}(q, e^{-t}) \\ &= q^{\kappa_{\mu^3/2}} e^{-|\mu^2|t/2} W_{\mu^2}(q, e^{-t}) \\ & \cdot \sum_{\eta} q^{(|\mu^1| - |\eta|)/2} s_{\mu^1/\eta}(E^{(\mu^2)^t}(u; q, e^{-t})) q^{(|\mu^3| - |\eta|)/2} s_{(\mu^3)^t/\eta}(E^{\mu^2}(u; q, e^{-t})). \end{aligned}$$

It is easy to see that the leading term of $W_{\mu^1, \mu^2, \mu^3}(q, e^{-t})$ is $\mathcal{W}_{\mu^1, \mu^2, \mu^3}(q)$. When $\mu^1 = (0)$,

$$\begin{aligned} W_{(0), \mu^2, \mu^3}(q, e^{-t}) &= q^{\kappa_{\mu^3/2}} e^{-|\mu^2|t/2} W_{\mu^2}(q, e^{-t}) q^{|\mu^3|/2} s_{(\mu^3)^t}(E^{\mu^2}(u; q, e^{-t})) \\ &= e^{-(|\mu^2| + |\mu^3|)t/2} q^{\kappa_{\mu^3/2}} W_{\mu^2, (\mu^3)^t}(q, e^{-t}). \end{aligned}$$

When $\mu^3 = (0)$,

$$W_{\mu^1, \mu^2, (0)}(q, e^{-t}) = e^{-|\mu^2|t/2} W_{\mu^2}(q, e^{-t}) q^{|\mu^1|/2} s_{\mu^1}(E^{(\mu^2)^t}(u; q, e^{-t})).$$

When $\mu^2 = (0)$,

$$(54) \quad \begin{aligned} & W_{\mu^1, (0), \mu^3}(q, e^{-t}) \\ &= q^{\kappa_{\mu^3/2}} \sum_{\eta} q^{(|\mu^1| - |\eta|)/2} s_{\mu^1/\eta}(E(u; q, e^{-t})) q^{(|\mu^3| - |\eta|)/2} s_{(\mu^3)^t/\eta}(E(u; q, e^{-t})). \end{aligned}$$

One then sees that W_{μ^1, μ^2, μ^3} does not have the $\mathbb{Z}_3$ cyclic symmetry.

Proposition 4.5. *We have*

$$(55) \quad W_{(\mu^1)^t, (\mu^2)^t, (\mu^3)^t}(q, e^{-t}) = (-1)^{|\mu^1| + |\mu^2| + |\mu^3|} W_{\mu^3, \mu^2, \mu^1}(q^{-1}, e^{-t}).$$

Proof. This is a straightforward consequence of (48). $\square$

5. FROM SUMMATIONS TO PRODUCTS

We use the results in the preceding two sections to prove some product expressions for certain sums of the invariants of the Hopf link and the deformed topological vertex. They generalize the results in [9, 10, 5, 6, 23] They will play a crucial role in the next section.

5.1. Results on $F_{\mu^1\mu^2}$. For two partitions μ^1, μ^2 , define:

$$\begin{aligned} F_{\mu^1\mu^2}(q, e^{-t}) &= \sum_{i \geq 1} (q^{\mu_i^1 - i + \frac{1}{2}} - e^{-t} q^{-i + \frac{1}{2}}) \sum_{j \geq 1} (q^{\mu_j^2 - j + \frac{1}{2}} - e^{-t} q^{-j + \frac{1}{2}}) \\ &\quad - \sum_{i \geq 1} (1 - e^{-t}) q^{-i + \frac{1}{2}} \sum_{j \geq 1} (1 - e^{-t}) q^{-j + \frac{1}{2}}. \end{aligned}$$

We have

$$(56) \quad F_{\mu^1\mu^2}(q, e^{-t}) = f_{\mu^1, \mu^2}(q) - e^{-t} f_{\mu^1}(q) - e^{-t} f_{\mu^2}(q).$$

Indeed, by (44) we have

$$\begin{aligned} &F_{\mu^1\mu^2}(q, e^{-t}) \\ &= qp_1(E^{\mu^1}(u; q, e^{-t}))p_1(E^{\mu^2}(u; q, e^{-t}) - qp_1(E^{(0)}(u; q, e^{-t}))p_1(E^{(0)}(u; q, e^{-t})) \\ &= q \left(\frac{1 - e^{-t}}{q - 1} + \sum_{i=1}^{l(\mu^1)} (q^{\mu_i^1 - i} - q^{-i}) \right) \left(\frac{1 - e^{-t}}{q - 1} + \sum_{j=1}^{l(\mu^2)} (q^{\mu_j^2 - j} - q^{-j}) \right) - \frac{(1 - e^{-t})^2 q}{(1 - q)^2} \\ &= (1 - e^{-t}) \frac{q}{q - 1} \sum_{i=1}^{l(\mu^1)} (q^{\mu_i^1 - i} - q^{-i}) + (1 - e^{-t}) \frac{q}{q - 1} \sum_{j=1}^{l(\mu^2)} (q^{\mu_j^2 - j} - q^{-j}) \\ &\quad + q \sum_{i=1}^{l(\mu^1)} (q^{\mu_i^1 - i} - q^{-i}) \cdot \sum_{j=1}^{l(\mu^2)} (q^{\mu_j^2 - j} - q^{-j}) \\ &= (1 - e^{-t}) f_{\mu^1}(q) + (1 - e^{-t}) f_{\mu^2}(q) + (q - 2 + q^{-1}) f_{\mu^1}(q) f_{\mu^2}(q) \\ &= f_{\mu^1\mu^2}(q) - e^{-t} f_{\mu^1}(q) - e^{-t} f_{\mu^2}(q). \end{aligned}$$

The expression

$$(57) \quad f_{\mu^1\mu^2} = \sum_{i,j=1}^{\infty} (q^{\mu_i^1 + \mu_j^2 - i - j + 1} - q^{-i - j + 1})$$

has been studied in [23]:

$$(58) \quad f_{\mu^1(\mu^2)^t}(q) = \sum_{(i,j) \in \mu^1} q^{\mu_i^1 + (\mu^2)_j^t - i - j + 1} + \sum_{(i,j) \in \mu^2} q^{-(\mu^1)_i^t - \mu_j^2 + i + j - 1}$$

$$(59) \quad = - \sum_{i,j \geq 1} (q^{\mu_i^1 - \mu_j^2 + j - i} - q^{j - i}).$$

In particular,

$$(60) \quad f_{\mu}(q) = \sum_{(i,j) \in \mu} q^{j - i}.$$

5.2. Some product expressions. By (56) we have

$$\begin{aligned} &\exp \left(\sum_{n \geq 1} \frac{u^n}{n} F_{\mu^1(\mu^2)^t}(q^n, e^{-nt}) \right) \\ &= \exp \left(\sum_{n \geq 1} \frac{u^n}{n} (f_{\mu^1(\mu^2)^t}(q^n) - e^{-nt} f_{\mu^1}(q^n) - e^{-nt} f_{(\mu^2)^t}(q^n)) \right). \end{aligned}$$

From this one can obtain various product expressions. First of all, by (57),

$$\begin{aligned}
 (61) \quad & \exp \left(\sum_{n \geq 1} \frac{u^n}{n} F_{\mu^1(\mu^2)^t}(q^n, e^{-nt}) \right) \\
 &= \prod_{i,j=1}^{\infty} \frac{(1 - uq^{-i-j+1})(1 - ue^{-t}q^{\mu_i^1-i-j+1})(1 - ue^{-t}q^{(\mu^2)_j^t-i-j+1})}{(1 - uq^{\mu_i^1+(\mu^2)_j^t-i-j+1})(1 - ue^{-t}q^{-i-j+1})(1 - ue^{-t}q^{-i-j+1})}.
 \end{aligned}$$

By (59),

$$\begin{aligned}
 (62) \quad & \exp \left(\sum_{n \geq 1} \frac{u^n}{n} F_{\mu^1(\mu^2)^t}(q^n, e^{-nt}) \right) \\
 &= \prod_{i,j=1}^{\infty} \frac{(1 - uq^{\mu_i^1-\mu_j^2-i+j})(1 - ue^{-t}q^{j-i})(1 - ue^{-t}q^{j-i})}{(1 - uq^{j-i})(1 - ue^{-t}q^{\mu_i^1-i+j})(1 - ue^{-t}q^{\mu_j^2-i+j})}.
 \end{aligned}$$

By (58) and (60),

$$\begin{aligned}
 (63) \quad & \exp \left(\sum_{n \geq 1} \frac{u^n}{n} F_{\mu^1(\mu^2)^t}(q^n, e^{-nt}) \right) \\
 &= \prod_{(i,j) \in \mu^1} \frac{1 - ue^{-t}q^{j-i}}{1 - uq^{\mu_i^1+(\mu^2)_j^t-i-j+1}} \cdot \prod_{(i,j) \in \mu^2} \frac{1 - ue^{-t}q^{-(j-i)}}{1 - uq^{-(\mu^1)_j^t+\mu_i^2-i-j+1}}.
 \end{aligned}$$

5.3. Product expressions for $\mathcal{K}_{\mu^1(\mu^2)^t}(Q)$. Define

$$\mathcal{K}_{\mu^1\mu^2}(Q; q, e^{-t}) = \sum_{\nu} Q^{|\nu|} e^{-(|\mu^1|+|\mu^2|+2|\nu|)t/2} W_{\mu^1\nu}(q, e^{-t}) W_{\nu\mu^2}(q, e^{-t}).$$

Proposition 5.1. *The following identities holds:*

$$\begin{aligned}
 (64) \quad & \frac{\mathcal{K}_{\mu^1\mu^2}(Q; q, e^{-t})}{\mathcal{K}_{(0)(0)}(Q; q, e^{-t})} \\
 &= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1}(q, e^{-t}) W_{\mu^2}(q, e^{-t}) \exp \left(\sum_{n=1}^{\infty} \frac{Q^n}{n} F_{\mu^1\mu^2}(q^n, e^{-nt}) \right).
 \end{aligned}$$

Proof. We have

$$\begin{aligned}
 & \mathcal{K}_{\mu^1\mu^2}(Q) \\
 &= \sum_{\nu} Q^{|\nu|} e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} q^{|\nu|/2} s_{\nu}(E^{\mu^1}(u; q, e^{-t})) W_{\mu^2} q^{|\nu|/2} s_{\nu}(E^{\mu^2}(u; q, e^{-t})) \\
 &= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} W_{\mu^2} \exp \sum_{n=1}^{\infty} \frac{(qQ)^n}{n} p_n(E^{\mu^1}(u; q, e^{-t})) p_n(E^{\mu^2}(u; q, e^{-t})).
 \end{aligned}$$

In particular, when $\mu^1 = \mu^2 = (0)$,

$$(65) \quad \mathcal{K}_{(0)(0)}(Q) = \exp \sum_{n=1}^{\infty} \frac{(qQ)^n}{n} p_n(E^{(0)}(u; q, e^{-t}))^2.$$

Hence

$$\begin{aligned}
& \frac{\mathcal{K}_{\mu^1\mu^2}(Q)}{\mathcal{K}_{(0)(0)}(Q)} \\
&= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} W_{\mu^2} \\
& \quad \exp \sum_{n=1}^{\infty} \frac{(qQ)^n}{n} (p_n(E^{\mu^1}(u; q, e^{-t})) p_n(E^{\mu^2}(u; q, e^{-t})) - p_n(E^{(0)}(u; q, e^{-t}))^2) \\
&= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} W_{\mu^2} \exp \left(\sum_{n \geq 1} \frac{(e^t Q)^n}{n} F_{\mu^1\mu^2}(q^n, e^{-nt}) \right).
\end{aligned}$$

□

We will find product expressions for $\mathcal{K}_{\mu^1(\nu^2)^t}$ in two different ways. By (65), (40) and Lemma 3.4 we have

$$(66) \quad \mathcal{K}_{(0)(0)} = \prod_{i,j=1}^{\infty} \frac{(1 - q^{-(i+j-1)}Q)^2}{(1 - q^{-(i+j-1)}e^tQ)(1 - q^{-(i+j-1)}e^{-t}Q)}$$

$$(67) \quad = \prod_{n=1}^{\infty} \frac{(1 - q^{-n}Q)^2}{(1 - q^{-n}e^tQ)(1 - q^{-n}e^{-t}Q)}.$$

Recall

$$(68) \quad W_{\mu}(q, e^{-t}) = \prod_{(i,j) \in \mu} \frac{e^{t/2}q^{(j-i)/2} - e^{-t/2}q^{-(j-i)/2}}{q^{(\mu_i+\mu_j^t-i-j+1)/2} - q^{-(\mu_i+\mu_j^t-i-j+1)/2}}.$$

It is possible to rewrite this as an infinite product. For the denominator, recall [23, Lemma 2.1],

$$(69) \quad \sum_{1 \leq i < j < \infty} (t^{\mu_i - \mu_j + j - i} - t^{j - i}) = - \sum_{(i,j) \in \mu} t^{\mu_i + \mu_j^t - i - j + 1},$$

for the numerator, recall

$$(70) \quad f_{\mu}(q) = \sum_{(i,j) \in \mu} q^{j-i} = - \sum_{i,j=1}^{\infty} (q^{\mu_i + j - i} - q^{j-i}).$$

Hence

$$\begin{aligned}
(71) \quad W_{\mu}(q, e^{-t}) &= \prod_{1 \leq i < j < \infty} \frac{q^{(\mu_i - \mu_j - i + j)/2} - q^{-(\mu_i - \mu_j - i + j)/2}}{q^{(j-i)/2} - q^{-(j-i)/2}} \\
&\quad \cdot \prod_{i,j=1}^{\infty} \frac{e^{t/2}q^{(\mu_i + j - i)/2} - e^{-t/2}q^{-(\mu_i + j - i)/2}}{e^{t/2}q^{(j-i)/2} - e^{-t/2}q^{-(j-i)/2}}.
\end{aligned}$$

By (62) and (63) one then gets two product expressions as follows.

Theorem 5.1. *We have*

$$\begin{aligned}
& \frac{\mathcal{K}_{\mu^1(\mu^2)^t}(Q)}{\mathcal{K}_{(0)(0)}(Q)} \\
&= e^{-(|\mu^1|+|\mu^2|)t/2} \prod_{(i,j) \in \mu^1} \frac{e^{t/2}q^{(j-i)/2} - e^{-t/2}q^{-(j-i)/2}}{q^{(\mu_i^1+(\mu^1)_j^t-i-j+1)/2} - q^{-(\mu_i^1+(\mu^1)_j^t-i-j+1)/2}} \\
(72) \quad & \cdot \prod_{(i,j) \in (\mu^2)^t} \frac{e^{t/2}q^{(j-i)/2} - e^{-t/2}q^{-(j-i)/2}}{q^{((\mu^2)_i^t+\mu_j^2-i-j+1)/2} - q^{-((\mu^2)_i^t+\mu_j^2-i-j+1)/2}} \\
& \cdot \prod_{(i,j) \in \mu^1} \frac{1 - e^{-t}Qq^{j-i}}{1 - Qq^{\mu_i^1+(\mu^2)_j^t-i-j+1}} \cdot \prod_{(i,j) \in \mu^2} \frac{1 - e^{-t}Qq^{-(j-i)}}{1 - Qq^{-(\mu^1)_j^t+\mu_i^2-i-j+1}}.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
& \frac{\mathcal{K}_{\mu^1(\mu^2)^t}(Q)}{\mathcal{K}_{(0)(0)}(Q)} \\
&= e^{-(|\mu^1|+|\mu^2|+2|\nu|)t/2} \prod_{1 \leq i < j < \infty} \frac{q^{(\mu_i^1-\mu_j^1-i+j)/2} - q^{-(\mu_i^1-\mu_j^1-i+j)/2}}{q^{(j-i)/2} - q^{-(j-i)/2}} \\
(73) \quad & \cdot \prod_{i,j=1}^{\infty} \frac{e^{t/2}q^{(\mu_i^1+j-i)/2} - e^{-t/2}q^{-(\mu_i^1+j-i)/2}}{e^{t/2}q^{(j-i)/2} - e^{-t/2}q^{-(j-i)/2}} \\
& \cdot \prod_{1 \leq i < j < \infty} \frac{q^{(\mu_i^2-\mu_j^2-i+j)/2} - q^{-(\mu_i^2-\mu_j^2-i+j)/2}}{q^{(j-i)/2} - q^{-(j-i)/2}} \\
& \cdot \prod_{i,j=1}^{\infty} \frac{e^{t/2}q^{(\mu_i^2+j-i)/2} - e^{-t/2}q^{-(\mu_i^2+j-i)/2}}{e^{t/2}q^{(j-i)/2} - e^{-t/2}q^{-(j-i)/2}} \\
& \cdot \prod_{i,j=1}^{\infty} \frac{(1 - uq^{\mu_i^1-\mu_j^2-i+j})(1 - ue^{-t}q^{j-i})(1 - ue^{-t}q^{j-i})}{(1 - uq^{j-i})(1 - ue^{-t}q^{\mu_i^1-i+j})(1 - ue^{-t}q^{\mu_j^2-i+j})}.
\end{aligned}$$

5.4. **Product expressions for $\mathcal{L}_{\mu^1(\mu^2)^t}(Q)$.** Define

$$\mathcal{L}_{\mu^1\mu^2}(Q; q, e^{-t}) = \sum_{\nu} e^{-(|\mu^1|+|\mu^2|+2|\nu|)t/2} (-Q)^{|\nu|} W_{\mu^1\nu}(q, e^{-t}) W_{\nu\mu^2}(q, e^{-t}).$$

Proposition 5.2. *The following identity holds:*

$$\begin{aligned}
& \frac{\mathcal{L}_{\mu^1\mu^2}(Q; q, e^{-t})}{\mathcal{L}_{(0)(0)}(Q; q, e^{-t})} \\
(74) \quad &= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1}(q, e^{-t}) W_{\mu^2}(q, e^{-t}) \exp \left(- \sum_{n=1}^{\infty} \frac{Q^n}{n} F_{\mu^1\mu^2}(q^n, e^{-nt}) \right).
\end{aligned}$$

Proof. We have

$$\begin{aligned}
& \mathcal{L}_{\mu^1 \mu^2}(Q) \\
&= \sum_{\nu} (-Q)^{|\nu|} e^{-|\mu^1|t/2} W_{\mu^1} q^{|\nu|/2} s_{\nu}(E^{\mu^1}(u; q, e^{-t})) \\
& \quad \cdot e^{-|\mu^2|t/2} W_{\mu^2} q^{|\nu|/2} s_{\nu^t}(E^{\mu^2}(u; q, e^{-t})) \\
&= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} W_{\mu^2} \exp \left(- \sum_{n=1}^{\infty} \frac{(qQ)^n}{n} p_n(E^{\mu^1}(u; q, e^{-t})) p_n(E^{\mu^2}(u; q, e^{-t})) \right).
\end{aligned}$$

In particular, when $\mu^1 = \mu^2 = (0)$,

$$(75) \quad \mathcal{L}_{(0)(0)}(Q) = \exp \left(- \sum_{n=1}^{\infty} \frac{(qQ)^n}{n} p_n(E^{(0)}(u; q, e^{-t}))^2 \right).$$

Hence

$$\begin{aligned}
& \frac{\mathcal{L}_{\mu^1 \mu^2}(Q)}{\mathcal{L}_{(0)(0)}(Q)} \\
&= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} W_{\mu^2} \exp \left(- \sum_{n=1}^{\infty} \frac{(qQ)^n}{n} (p_n(E^{\mu^1}(u; q, e^{-t})) p_n(E^{\mu^2}(u; q, e^{-t})) \right. \\
& \quad \left. - p_n(E^{(0)}(u; q, e^{-t}))^2) \right) \\
&= e^{-(|\mu^1|+|\mu^2|)t/2} W_{\mu^1} W_{\mu^2} \exp \left(- \sum_{n \geq 1} \frac{Q^n}{n} F_{\mu^1 \mu^2}(q^n, e^{-nt}) \right).
\end{aligned}$$

□

By (62) and (63) one then gets two product expressions.

Theorem 5.2.

$$\begin{aligned}
& \frac{\mathcal{L}_{\mu^1(\mu^2)^t}(Q)}{\mathcal{L}_{(0)(0)}(Q)} \\
&= \prod_{(i,j) \in \mu^1} \frac{e^{t/2} q^{(j-i)/2} - e^{-t/2} q^{-(j-i)/2}}{q^{(\mu_i^1 + (\mu^1)_j^t - i - j + 1)/2} - q^{-(\mu_i^1 + (\mu^1)_j^t - i - j + 1)/2}} \\
& \quad \cdot \prod_{(i,j) \in (\mu^2)^t} \frac{e^{t/2} q^{(j-i)/2} - e^{-t/2} q^{-(j-i)/2}}{q^{((\mu^2)_i^t + \mu_j^2 - i - j + 1)/2} - q^{-((\mu^2)_i^t + \mu_j^2 - i - j + 1)/2}} \\
& \quad \cdot \prod_{(i,j) \in \mu^1} \frac{1 - Q q^{\mu_i^1 + (\mu^2)_j^t - i - j + 1}}{1 - Q e^{-t} q^{j-i}} \cdot \prod_{(i,j) \in \mu^2} \frac{1 - Q q^{-(\mu^1)_j^t + \mu_i^2 - i - j + 1}}{1 - Q e^{-t} q^{-(j-i)}}.
\end{aligned}$$

In particular, when $\mu^1 = \mu^2 = \mu$,

$$\begin{aligned}
& \frac{\mathcal{L}_{\mu \mu^t}(Q)}{\mathcal{L}_{(0)(0)}(Q)} \\
(76) \quad &= (-1)^{|\mu|} \prod_{e \in \mu} \frac{(1 - e^{-t} q^{c(e)})(1 - e^{-t} q^{-c(e)})}{(1 - Q e^{-t} q^{c(e)})(1 - Q e^{-t} q^{-c(e)})} \cdot \frac{(1 - Q q^{h(e)})(1 - Q q^{-h(e)})}{(1 - q^{h(e)})(1 - q^{-h(e)})}.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
& \frac{\mathcal{L}_{\mu^1(\mu^2)^t}(Q)}{\mathcal{L}_{(0)(0)}(Q)} \\
&= \prod_{1 \leq i < j < \infty} \frac{q^{(\mu_i^1 - \mu_j^1 - i + j)/2} - q^{-(\mu_i^1 - \mu_j^1 - i + j)/2}}{q^{(j-i)/2} - q^{-(j-i)/2}} \\
&\quad \cdot \prod_{i,j=1}^{\infty} \frac{e^{t/2} q^{(\mu_i^1 + j - i)/2} - e^{-t/2} q^{-(\mu_i^1 + j - i)/2}}{e^{t/2} q^{(j-i)/2} - e^{-t/2} q^{-(j-i)/2}} \\
&\quad \cdot \prod_{1 \leq i < j < \infty} \frac{q^{(\mu_i^2 - \mu_j^2 - i + j)/2} - q^{-(\mu_i^2 - \mu_j^2 - i + j)/2}}{q^{(j-i)/2} - q^{-(j-i)/2}} \\
&\quad \cdot \prod_{i,j=1}^{\infty} \frac{e^{t/2} q^{(\mu_i^2 + j - i)/2} - e^{-t/2} q^{-(\mu_i^2 + j - i)/2}}{e^{t/2} q^{(j-i)/2} - e^{-t/2} q^{-(j-i)/2}} \\
&\quad \cdot \prod_{i,j=1}^{\infty} \frac{(1 - uq^{\mu_i^1 - \mu_j^2 - i + j})(1 - ue^{-t} q^{j-i})(1 - ue^{-t} q^{j-i})}{(1 - uq^{j-i})(1 - ue^{-t} q^{\mu_i^1 - i + j})(1 - ue^{-t} q^{\mu_j^2 - i + j})}.
\end{aligned} \tag{77}$$

5.5. **Generalizations.** Define:

$$\tilde{\mathcal{K}}_{\mu^1 \dots \mu^N}(Q_1, \dots, Q_{N-1}; q, e^{-t}) = \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N q^{\kappa_{\nu^k}/2} W_{\nu^{k-1} \mu^k(\nu^k)^t}(q, e^{-t}) Q_k^{|\nu^k|},$$

where $\nu^0 = \nu^N = (0)$, $Q_N = 1$.

Proposition 5.3. *We have the following identity:*

$$\begin{aligned}
& \frac{\tilde{\mathcal{K}}_{\mu^1 \dots \mu^N}(Q_1, \dots, Q_{N-1})}{\prod_{1 \leq k < l \leq N} \mathcal{K}_{(0)(0)}(Q_k \cdots Q_{l-1})} \\
&= \prod_{k=1}^N W_{\mu^k}(q, e^{-t}) \cdot \exp \left(\sum_{1 \leq k < l \leq N} \sum_{n \geq 1} \frac{(Q_k \cdots Q_{l-1})^n}{n} F_{\mu^i(\mu^j)^t}(q^n, e^{-nt}) \right).
\end{aligned}$$

Proof.

$$\begin{aligned}
& \tilde{\mathcal{K}}_{\mu^1 \dots \mu^N}(Q_1, \dots, Q_{N-1}; q, e^{-t}) \\
&= \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N q^{\kappa_{\nu^k}/2} W_{\nu^{i-1} \mu^k (\nu^k)^t}(q, e^{-t}) Q_k^{|\nu^k|} \quad (\nu^0 = \nu^N = (0)) \\
&= \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N e^{-|\mu^k|t/2} W_{\mu^k}(q, e^{-t}) \\
&\quad \cdot \sum_{\eta^{k-1}} q^{(|\nu^{k-1}| - |\eta^{k-1}|)/2} s_{\nu^{k-1}/\eta^{k-1}}(E^{(\mu^k)^t} q, e^{-t}) \\
&\quad \cdot q^{(|\nu^k| - |\eta^{k-1}|)/2} s_{\nu^k/\eta^{k-1}}(E^{\mu^k}(q, e^{-t})) Q_k^{|\nu^k|} \\
&\quad \quad \quad (\nu^0 = \nu^N = \eta^0 = \eta^{N-1} = (0)) \\
&= \prod_{k=1}^N e^{-|\mu^k|t/2} W_{\mu^k}(q, e^{-t}) \sum_{\nu^1, \dots, \nu^{N-1}} \sum_{\eta^1, \dots, \eta^{N-2}} \prod_{k=1}^{N-1} q^{(|\nu^k| - |\eta^{k-1}|)/2} \\
&\quad \cdot s_{\nu^k/\eta^{k-1}}(E^{\mu^k}(q, e^{-t})) q^{(|\nu^k| - |\eta^k|)/2} Q_k^{|\nu^k|} s_{\nu^k/\eta^k}(E^{(\mu^{k+1})^t}(q, e^{-t})) \\
&\quad \quad \quad (\eta^0 = \eta^{N-1} = (0)) \\
&= \prod_{k=1}^N e^{-|\mu^k|t/2} W_{\mu^k}(q, e^{-t}) \cdot \prod_{1 \leq k < l \leq N} \exp \sum_{n=1}^{\infty} \frac{(q Q_k Q_{k+1} \dots Q_l)^n}{n} \\
&\quad \cdot p_n(E^{\mu^k}(q, e^{-t})) p_n(E^{(\mu^l)^t}(q, e^{-t})).
\end{aligned}$$

In the last equality we have used (21). Hence

$$\begin{aligned}
& \frac{\tilde{\mathcal{K}}_{\mu^1 \dots \mu^N}(Q_1, \dots, Q_{N-1}; q, e^{-t})}{\prod_{1 \leq k < l \leq N} \mathcal{K}_{(0)(0)}(Q_k \dots Q_{l-1}; q, e^{-t})} \\
&= \prod_{k=1}^N (e^{-|\mu^k|t/2} W_{\mu^k}(q, e^{-t})) \cdot \exp \sum_{1 \leq k < l \leq N} \sum_{n \geq 1} \frac{(Q_k \dots Q_{l-1})^n}{n} F_{\mu^k(\mu^l)^t}(q^n, e^{-nt}).
\end{aligned}$$

□

Hence one can easily get a Nekrasov type infinite product expression and the following product expression:

$$\begin{aligned}
& \frac{\tilde{\mathcal{K}}_{\mu^1 \dots \mu^N}(Q_1, \dots, Q_{N-1}; q, e^{-t})}{\prod_{1 \leq k < l \leq N} \mathcal{K}_{(0)(0)}(Q_k \dots Q_{l-1}; q, e^{-t})} \\
&= \prod_{\alpha=1}^N (e^{-|\mu^\alpha|t/2} \prod_{(i,j) \in \mu^\alpha} \frac{e^{t/2} q^{(j-i)/2} - e^{-t/2} q^{-(j-i)/2}}{q^{h(i,j)/2} - q^{-h(i,j)/2}}) \\
&\quad \cdot \prod_{1 \leq \alpha < \gamma \leq N} \left(\prod_{(i,j) \in \mu^\alpha} \frac{1 - Q_\alpha \dots Q_{\gamma-1} e^{-t} q^{j-i}}{1 - Q_\alpha \dots Q_{\gamma-1} q^{\mu_i^\alpha + (\mu^\gamma)_j^t - i + j + 1}} \right. \\
&\quad \cdot \left. \prod_{(i,j) \in \mu^\gamma} \frac{1 - Q_\alpha \dots Q_{\gamma-1} e^{-t} q^{-(j-i)}}{1 - Q_\alpha \dots Q_{\gamma-1} q^{\mu_j^\gamma + (\mu^\alpha)_i^t - i + j + 1}} \right)
\end{aligned} \tag{78}$$

6. DEFORMED TOPOLOGICAL STRING PARTITION FUNCTIONS OF SOME LOCAL CALABI-YAU GEOMETRIES

In this section we introduce some deformed topological string partition functions for some local Calabi-Yau geometries. We identify them with the deformed instanton counting partition functions considered in §2. Our results generalize those related to Nekrasov's partition function [18, 9, 5, 23].

6.1. The resolved conifold case. The web diagram is




Figure 1

The (undeformed) partition function is given by:

$$Z_{(0)} = \sum_{\mu} \mathcal{W}_{\mu^t}(q) (-1)^{|\mu|} Q^{|\mu|} \mathcal{W}_{\mu}(q).$$

We take the deformed partition function to be:

$$(79) \quad \mathcal{Z}_{(0)} = \sum_{\mu} e^{-|\mu|t} W_{\mu^t}(q, e^{-t}) (-1)^{|\mu|} Q^{|\mu|} W_{\mu}(q, e^{-t}).$$

By (30) it can be also written as:

$$(80) \quad \mathcal{Z}_{(0)} = \sum_{\mu} e^{-|\mu|t} Q^{|\mu|} W_{\mu}(q, e^{-t}) W_{\mu^t}(q, e^t).$$

By (28), one easily gets:

$$(81) \quad \mathcal{Z}_{(0)} = \sum_{\mu} \prod_{(i,j) \in \mu} \frac{(1 - e^{-t} q^{j-i})(1 - e^{-t} q^{i-j})}{(1 - q^{h(i,j)})(1 - q^{-h(i,j)})}.$$

We will prove two results for such deformed partition functions. First of all, we will identify it with certain partition functions on the Hilbert schemes. Secondly, we will find infinite product expression for it so that one can extract the *deformed Gopakumar-Vafa invariants*.

By comparing (81) and (5), one gets the following result which generalizes the undeformed case considered in [7, 12]:

Theorem 6.1. *One has the following identification of deformed partition functions:*

$$(82) \quad \mathcal{Z}_{(0)} = \sum_{n=0}^{\infty} Q^n \chi((\mathbb{C}^2)^{[n]}, \Lambda_{-e^{-t}}(V_n) \otimes \Lambda_{-e^{-t}}(V_n^*))(q, q^{-1}).$$

The second result is a straightforward consequence of (75) and the observation

$$\mathcal{Z}_{(0)} = \mathcal{L}_{(0)(0)}(Q).$$

It can be stated as follows.

Theorem 6.2. *The deformed partition function for the resolved conifold can be rewritten as follows.*

$$(83) \quad \mathcal{Z}_{(0)} = \exp\left(-\sum_{n=1}^{\infty} \frac{(1-e^{-nt})^2}{(q^{n/2}-q^{-n/2})^2} Q^n\right)$$

$$(84) \quad = \prod_{n=1}^{\infty} \frac{(1-Qq^n)^n (1-Qe^{-t}q^n)^n}{(1-Qe^{-t/2}q^n)^{2n}}.$$

Besides the natural T^2 -action on $\mathbb{C}^2$, consider an extra circle action on V_1 and V_1^* by scalar multiplications along the fibers. Then we have

$$\chi(\mathbb{C}^2, \Lambda_{-1}(V_1) \otimes \Lambda_{-1}(V_1^*))(t_1, t_2, e^{-t}) = \frac{(1-e^{-t})^2}{(1-t_1^{-1})(1-t_2^{-1})}.$$

Then we have

$$(85) \quad \mathcal{Z}_{(0)} = \exp \sum_{n=1}^{\infty} \frac{1}{n} \chi(\mathbb{C}^2, \Lambda_{-1}(V_1) \otimes \Lambda_{-1}(V_1^*))(t_1^n, t_2^n, e^{-nt})|_{t_1=q, t_2=q^{-1}}.$$

From this one expects an interpretation in terms of symmetric products generalizing the nondeformed case in [12].

6.2. A related case. A related case is given by the following diagram, where the two horizontal edges are glued with each other.




Figure 2

The (undeformed) partition function is given by

$$(86) \quad Z(Q, Q_m, q) = \sum_{\mu, \nu} (-Q)^{|\mu|} (-Q_m)^{|\nu|} \mathcal{W}_{\mu\nu}(q) \mathcal{W}_{\nu^t \mu^t}(q).$$

We take the deformed partition function to be

$$(87) \quad \tilde{Z}(Q, Q_m, q, e^{-t}) = \sum_{\mu, \nu} (-Q)^{|\mu|} (-Q_m)^{|\nu|} W_{\mu\nu}(q, e^{-t}) W_{\nu^t \mu^t}(q, e^{-t}).$$

Now note that

$$(88) \quad \tilde{Z}(Q, Q_m, q, e^{-t}) = \sum_{\mu} (-Q)^{|\mu|} \mathcal{L}_{\mu(\mu)^t}(Q_m).$$

Hence by (76)

$$\begin{aligned}
(89) \quad & \frac{\tilde{Z}(Q, Q_m, q, e^{-t})}{Z_{(0)}(Q_m, q, e^{-t})} \\
&= \sum_{\mu} Q^{|\mu|} \prod_{e \in \mu} \frac{(1 - e^{-t} q^{c(e)})(1 - e^{-t} q^{-c(e)})}{(1 - Q_m e^{-t} q^{c(e)})(1 - Q_m e^{-t} q^{-c(e)})} \\
&\quad \cdot \prod_{e \in \mu} \frac{(1 - Q_m q^{h(e)})(1 - Q_m q^{-h(e)})}{(1 - q^{h(e)})(1 - q^{-h(e)})}.
\end{aligned}$$

We now identify this as a partition function on the Hilbert schemes by comparing with (8), generalizing the results in the deformed case [7, 12]:

Theorem 6.3. *One has the following identification of deformed partition functions:*

$$(90) \quad \frac{\tilde{Z}(Q, Q_m)}{Z_{(0)}(Q_m)} = \sum_{n=0}^{\infty} Q^n \chi((\mathbb{C}^2)^{[n]}, \Lambda_{-Q_m}(\mathbb{V}_n) \otimes W_{e^{-t}, Q_m}(\mathbb{V}_n^*))(q, q^{-1}).$$

We do not know a rigorous method to establish an infinite product expression for $\tilde{Z}(Q, Q_m)$ at present. For the undeformed partition function, a physical calculation based on duality with Chern-Simons theory was carried out in [7], and a mathematical proof based on symmetric products was given in [12]. Here we first present another mathematical proof for the undeformed case using Schur calculus.

Theorem 6.4. [7, 12] *For the undeformed partition function, one has*

$$(91) \quad Z(Q, Q_m, q) = \prod_{k=0}^{\infty} \prod_{l=1}^{\infty} \frac{(1 - Q^{k+1} Q_m^k q^l)^l (1 - Q^k Q_m^{k+1} q^l)^l}{(1 - Q^{k+1} Q_m^{k+1} q^l)^{2l}}.$$

Proof. We use the following identity [21]:

$$(92) \quad W_{\mu\nu}(q) = (-1)^{|\mu|+|\nu|} q^{(\kappa_{\mu}+\kappa_{\nu})/2} \sum_{\eta} s_{\mu/\eta}(q^{-\rho}) s_{\nu/\rho}(q^{-\rho}).$$

and (25):

$$\begin{aligned}
Z(Q, Q_m, q) &= \sum_{\mu, \nu} (-Q)^{|\mu|} (-Q_m)^{|\nu|} W_{\mu\nu}(q) W_{\nu^t \mu^t}(q) \\
&= \sum_{\mu, \nu, \xi, \eta} (-Q)^{|\mu|} (-Q_m)^{|\nu|} s_{\mu/\xi}(q^{-\rho}) s_{\nu/\xi}(q^{-\rho}) s_{\nu^t/\eta}(q^{-\rho}) s_{\mu^t/\eta}(q^{-\rho}) \\
&= \sum_{\mu, \nu, \xi, \eta} Q^{|\mu|} Q_m^{|\nu|} s_{\mu/\xi}(q^{-\rho}) s_{\nu/\xi}(q^{-\rho}) s_{\nu/\eta}(q^{\rho}) s_{\mu/\eta}(q^{\rho}) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n(1 - (QQ_m)^n)} (Q^n p_n(q^{-\rho}) p_n(q^{\rho}) + Q_m^n p_n(q^{-\rho}) p_n(q^{\rho}) \\
&\quad + (QQ_m)^n p_n(q^{-\rho}) p_n(q^{-\rho}) + (QQ_m)^n p_n(q^{\rho}) p_n(q^{\rho})).
\end{aligned}$$

By (26) we then have:

$$\begin{aligned}
& Z(Q, Q_m, q) \\
&= \exp \sum_{n=1}^{\infty} \frac{1}{n(1 - (QQ_m)^n)} \left(-\frac{Q^n q^n}{(1 - q^n)^2} - \frac{Q_m^n q^n}{(1 - q^n)^2} + \frac{2(QQ_m)^n q^n}{(1 - q^n)^2} \right) \\
&= \prod_{k=0}^{\infty} \prod_{l=1}^{\infty} \frac{(1 - Q^{k+1} Q_m^k q^l)^l (1 - Q^k Q_m^{k+1} q^l)^l}{(1 - Q^{k+1} Q_m^{k+1} q^l)^{2l}}.
\end{aligned}$$

□

It turns out that this proof can be generalized to another deformed partition function:

$$(93) \quad \mathcal{Z}(Q, Q_m; q, e^{-t}) = \sum_{\nu, \eta} (-Q_1)^{|\nu|} (-Q_m)^{|\eta|} W_{\eta(0)\nu}(q, e^{-t}) W_{\eta^t(0)\nu^t}(q, e^{-t}).$$

Theorem 6.5. *We have*

$$\begin{aligned}
& \mathcal{Z}(Q, Q_m; q, e^{-t}) \\
&= \prod_{k=0}^{\infty} \left(\prod_{l=1}^{\infty} \left(\frac{(1 - Q^{k+1} Q_m^k q^l)(1 - e^{-2t} Q^{k+1} Q_m^k q^l)}{(1 - e^{-t} Q^{k+1} Q_m^k q^l)^2} \right)^l \right. \\
(94) \quad & \cdot \left(\frac{(1 - Q^k Q_m^{k+1} q^l)(1 - e^{-2t} Q^k Q_m^{k+1} q^l)}{(1 - e^{-t} Q^k Q_m^{k+1} q^l)^2} \right)^l \\
& \cdot \left. \left(\frac{(1 - e^{-t} Q^{k+1} Q_m^{k+1} q^l)^2}{(1 - Q^{k+1} Q_m^{k+1} q^l)(1 - e^{-2t} Q^{k+1} Q_m^{k+1} q^l)} \right)^{2l} \right).
\end{aligned}$$

Proof. By (25) we have:


$$\begin{aligned}
& \mathcal{Z}(Q, Q_m; q, e^{-t}) \\
&= \sum_{\nu, \eta} (-Q)^{|\nu|} (-Q_m)^{|\eta|} \\
& \cdot \sum_{\xi^1} q^{(|\eta| - |\xi^1|)/2} s_{\eta/\xi^1}(E(u; q, e^{-t})) q^{(|\nu| - |\xi^1|)/2} s_{\nu^t/\xi^1}(E(u; q, e^{-t})) \\
& \cdot \sum_{\xi^2} q^{(|\eta^t| - |\xi^2|)/2} s_{\eta^t/\xi^2}(E(u; q, e^{-t})) q^{(|\nu| - |\xi^2|)/2} s_{\nu/\xi^2}(E(u; q, e^{-t})) \\
&= \exp \sum_{n=1}^{\infty} \frac{q^n}{n(1 - (Q_1 Q_m)^n)} \\
& (-Q^n - Q_m^n + 2(QQ_m)^n) p_n(E(u; q, e^{-t})) p_n(E(u; q, e^{-t})) \\
&= \exp \sum_{n=1}^{\infty} \frac{q^n}{n(1 - (Q_1 Q_m)^n)} \frac{(1 - e^{-nt})^2}{(q^n - 1)^2} (-Q^n - Q_m^n + 2(QQ_m)^n)
\end{aligned}$$

It is then an easy exercise to show that:

$$\begin{aligned} \mathcal{Z}(Q, Q_m; q, e^{-t}) &= \prod_{k=0}^{\infty} \left(\prod_{l=1}^{\infty} \left(\frac{(1 - Q^{k+1} Q_m^k q^l)(1 - e^{-2t} Q^{k+1} Q_m^k q^l)}{(1 - e^{-t} Q^{k+1} Q_m^k q^l)^2} \right)^l \right. \\ &\quad \cdot \left(\frac{(1 - Q^k Q_m^{k+1} q^l)(1 - e^{-2t} Q^k Q_m^{k+1} q^l)}{(1 - e^{-t} Q^k Q_m^{k+1} q^l)^2} \right)^l \\ &\quad \cdot \left. \left(\frac{(1 - e^{-t} Q^{k+1} Q_m^{k+1} q^l)^2}{(1 - Q^{k+1} Q_m^{k+1} q^l)(1 - e^{-2t} Q^{k+1} Q_m^{k+1} q^l)} \right)^{2l} \right). \end{aligned}$$

□

6.3. Another related case. Consider the local Calabi-Yau geometry with the following web diagram:



where we glue together the horizontal edge with other and also the vertex edge with each other. The undeformed theory of this case has been discussed in Section 5.1.1 and Section 6.2 in [7]. The partition function by topological vertex method is

$$(95) \quad Z(Q, Q_m, Q_1; q) = \sum_{\mu, \nu, \eta} (-Q)^{|\mu|} (-Q_1)^{|\nu|} (-Q_m)^{|\eta|} \mathcal{W}_{\mu\nu\eta}(q) \mathcal{W}_{\mu^t \nu^t \eta^t}(q).$$

We take the deformed partition function to be:

$$(96) \quad \begin{aligned} &\mathcal{Z}(Q, Q_m, Q_1; q, e^{-t}) \\ &= \sum_{\mu, \nu, \eta} (-Q)^{|\mu|} (-Q_1)^{|\nu|} (-Q_m)^{|\eta|} W_{\eta\mu\nu}(q, e^{-t}) W_{\eta^t \mu^t \nu^t}(q, e^{-t}). \end{aligned}$$

Theorem 6.6. *We have*

$$(97) \quad \begin{aligned} &\frac{\mathcal{Z}(Q, Q_m, Q_1; q, e^{-t})}{\tilde{\mathcal{Z}}(Q_1, Q_m; q, e^{-t})} \\ &= \sum_{\mu} Q^{|\mu|} \prod_{(i,j) \in \mu} \frac{(1 - e^{-t} q^{j-i})(1 - e^{-t} q^{-(j-i)})}{(1 - q^{h(i,j)})(1 - q^{-h(i,j)})} \\ &\quad \prod_{k=1}^{\infty} \frac{(1 - Q_m Q_{\rho}^{k-1} q^{h(i,j)})(1 - Q_m Q_{\rho}^{k-1} q^{-h(i,j)})}{(1 - e^{-t} Q_m Q_{\rho}^{k-1} q^{j-i})(1 - e^{-t} Q_m Q_{\rho}^{k-1} q^{-(j-i)})} \\ &\quad \frac{(1 - Q_1 Q_{\rho}^{k-1} q^{h(i,j)})(1 - Q_1 Q_{\rho}^{k-1} q^{-h(i,j)})}{(1 - e^{-t} Q_1 Q_{\rho}^{k-1} q^{j-i})(1 - e^{-t} Q_1 Q_{\rho}^{k-1} q^{-(j-i)})} \\ &\quad \frac{(1 - Q_{\rho}^k e^{-t} q^{j-i})^2 (1 - Q_{\rho}^k e^{-t} q^{-(j-i)})^2}{(1 - Q_{\rho}^k q^{h(i,j)})^2 (1 - Q_{\rho}^k q^{-h(i,j)})^2}. \end{aligned}$$

Proof. First of all, by the definition of the deformed topological vertex we have:

$$\begin{aligned}
& \mathcal{Z}(Q, Q_m, Q_1; q, e^{-t}) \\
&= \sum_{\mu, \nu, \eta} (-Q)^{|\mu|} (-Q_1)^{|\nu|} (-Q_m)^{|\eta|} e^{-|\mu|t/2} W_\mu(q, e^{-t}) e^{-|\mu^t|t/2} W_{\mu^t}(q, e^{-t}) \\
&\quad \cdot \sum_{\xi^1} q^{(|\eta| - |\xi^1|)/2} s_{\eta/\xi^1}(E^{\mu^t}(u; q, e^{-t})) q^{(|\nu| - |\xi^1|)/2} s_{\nu^t/\xi^1}(E^\mu(u; q, e^{-t})) \\
&\quad \cdot \sum_{\xi^2} q^{(|\eta^t| - |\xi^2|)/2} s_{\eta^t/\xi^2}(E^\mu(u; q, e^{-t})) q^{(|\nu| - |\xi^2|)/2} s_{\nu/\xi^2}(E^{\mu^t}(u; q, e^{-t})).
\end{aligned}$$

Hence by (25) we have:

$$\begin{aligned}
& \mathcal{Z}(Q, Q_m, Q_1; q, e^{-t}) \\
&= e^{-|\mu|t} W_\mu(q, e^{-t}) W_{\mu^t}(q, e^{-t}) \exp \sum_{n=1}^{\infty} \frac{q^n}{n(1 - (Q_1 Q_m)^n)} \\
&\quad \cdot (-Q_1^n - Q_m^n + 2(Q_1 Q_m)^n) p_n(E^\mu(u; q, e^{-t})) p_n(E^{\mu^t}(u; q, e^{-t})).
\end{aligned}$$

It follows that

$$\begin{aligned}
& \frac{\mathcal{Z}(Q, Q_m, Q_1; q, e^{-t})}{\mathcal{Z}(Q_1, Q_m; q, e^{-t})} \\
&= e^{-|\mu|t} W_\mu(q, e^{-t}) W_{\mu^t}(q, e^{-t}) \exp \sum_{n=1}^{\infty} \frac{q^n}{n(1 - (Q_1 Q_m)^n)} \\
&\quad \cdot (-Q_1^n - Q_m^n + 2(Q_1 Q_m)^n) F_{\mu, \mu^t}(q^n, e^{-nt}).
\end{aligned}$$

Hence (97) follows from (63). $\square$

Theorem 6.7. *We have the following identity:*

$$\begin{aligned}
(98) \quad & \frac{\mathcal{Z}(Q, Q_m, Q_1; q, e^{-t})}{\mathcal{Z}(Q_1, Q_m; q, e^{-t})} \\
&= \sum_{n \geq 0} Q^n \chi((\mathbb{C}^2)^{[n]}, \text{Ell}(T(\mathbb{C}^2)^{[n]}, \mathbb{V}_n)(Q_m, Q_\rho, e^{-t}))(q, q^{-1}),
\end{aligned}$$

where $Q_\rho = Q_1 Q_m$.

Proof. we rewrite the right-hand side of (97) as follows.

$$\begin{aligned}
& \sum_{\mu} Q^{|\mu|} \prod_{(i,j) \in \mu} \prod_{k=1}^{\infty} \left(\frac{(1 - Q_\rho^{k-1} Q_m q^{h(i,j)})(1 - Q_\rho^{k-1} Q_m q^{-h(i,j)})}{(1 - Q_\rho^{k-1} q^{h(i,j)})(1 - Q_\rho^{k-1} q^{-h(i,j)})} \right. \\
&\quad \cdot \frac{(1 - Q_\rho^k Q_m^{-1} q^{h(i,j)})(1 - Q_\rho^k Q_m^{-1} q^{-h(i,j)})}{(1 - Q_\rho^k q^{h(i,j)})(1 - Q_\rho^k q^{-h(i,j)})} \\
&\quad \cdot \frac{(1 - e^{-t} Q_\rho^{k-1} q^{c(i,j)})(1 - e^{-t} Q_\rho^{k-1} q^{-c(i,j)})}{(1 - e^{-t} Q_\rho^{k-1} Q_m q^{c(i,j)})(1 - e^{-t} Q_\rho^{k-1} Q_m q^{-c(i,j)})} \\
&\quad \cdot \left. \frac{(1 - e^{-t} Q_\rho^k q^{c(i,j)})(1 - e^{-t} Q_\rho^k q^{-c(i,j)})}{(1 - e^{-t} Q_\rho^k Q_m^{-1} q^{c(i,j)})(1 - e^{-t} Q_\rho^k Q_m^{-1} q^{-c(i,j)})} \right),
\end{aligned}$$

Then (98) follows from (10). $\square$

6.4. **The $SU(2)$ case.** The web diagrams are:




Figure 3

where $B, F \in H_2(\mathbb{F}_m, \mathbb{Z})$ are the homological classes of the base and the fiber respectively, and we have

$$F^2 = 0, \quad B^2 = -m, \quad BF = 1,$$

hence

$$(B + mF)^2 = m.$$

The undeformed partition functions are calculated in physics literature in [2] and formulated as a Feynman rule in [8]. The mathematical proof can be found in [22]. The self-intersection numbers of the divisors in the surfaces represented by the internal edges play a role in the formula:

$$\begin{aligned} Z_{\mathbb{F}_m}(Q_B, Q_F) &= \sum_{\mu^{1,2}, \nu^{1,2}} \mathcal{W}_{\mu^1 \nu^1}(q) Q_F^{|\nu^1|} \mathcal{W}_{\nu^1 (\mu^2)^t}(q) (-1)^{m|\mu^2|} q^{m\kappa_{(\mu^2)^t}} \\ &\quad \cdot (Q_B Q_F^m)^{|\mu^2|} \mathcal{W}_{(\mu^2)^t \nu^2}(q) Q_F^{|\nu^2|} \mathcal{W}_{\nu^2 \mu^1}(q) (-1)^{-m|\mu^1|} q^{-m\kappa_{\mu^1}} \\ &= \sum_{\mu^1, \mu^2} Q_B^{|\mu^1| + |\mu^2|} [(-1)^{|\mu^1| + |\mu^2|} Q_F^{|\mu^2|} q^{-\frac{1}{2}(\kappa_{\mu^1} + \kappa_{\mu^2})}]^m K_{\mu^1, (\mu^2)^t}(Q_F)^2. \end{aligned}$$

Here the locations of the partitions are indicated below:




Figure 4

The above expression for $Z_{\mathbb{F}_m}$ suggests the following partition function:

$$\begin{aligned} &\tilde{Z}_{\mathbb{F}_m}(q, Q_B, Q_F, e^{-t}) \\ &= \sum_{\mu^{1,2}, \nu^{1,2}} e^{-(2|\mu^1| + 2|\mu^2| + |\nu^1| + |\nu^2|)t} \\ (99) \quad &\cdot W_{\mu^1 \nu^1}(q, e^{-t}) Q_F^{|\nu^1|} W_{\nu^1 (\mu^2)^t}(q, e^{-t}) (-1)^{m|\mu^2|} q^{m\kappa_{(\mu^2)^t}} \\ &\cdot (Q_B Q_F^m)^{|\mu^2|} W_{(\mu^2)^t \nu^2}(q, e^t) Q_F^{|\nu^2|} W_{\nu^2 \mu^1}(q, e^t) (-1)^{-m|\mu^1|} q^{-m\kappa_{\mu^1}}. \end{aligned}$$

This is not a deformation of $Z_{\mathbb{F}_m}$. However it has a similar property as $Z_{\mathbb{F}_m}$ as follows.

Theorem 6.8. *We have*

$$(100) \quad \frac{\mathcal{Z}_{\mathbb{F}_m}(Q_B, Q_F; q, e^{-t})}{\mathcal{K}_{(0)(0)}(Q_F; q, e^{-t})\mathcal{K}_{(0)(0)}(Q_F; q, e^t)} = \sum_{k \geq 0} Q_B^k \chi(M(2, k), \Lambda_{-e^{-t}}(\mathbb{V}_k \otimes \mathbb{W}_k^*) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_k^* \otimes \mathbb{W}_k) \otimes E_{N,k}^m)(e_1, e_2, q, q^{-1})$$

where $Q_F = e_1 e_2^{-1}$, $e_1 = -1$.

Proof. By (99) we have

$$\begin{aligned} & \frac{\mathcal{Z}_{\mathbb{F}_m}(q, Q_B, Q_F, e^{-t})}{\mathcal{K}_{(0)(0)}(Q_F; q, e^{-t})\mathcal{K}_{(0)(0)}(Q_F; q, e^t)} \\ &= \sum_{\mu^1, \mu^2} Q_B^{|\mu^1|+|\mu^2|} [(-1)^{|\mu^1|+|\mu^2|} Q_F^{|\mu^2|} q^{-\frac{1}{2}(\kappa_{\mu^1} + \kappa_{\mu^2})}]^m \\ & \quad \cdot \frac{\mathcal{K}_{\mu^1, (\mu^2)^t}(Q_F; q, e^{-t})\mathcal{K}_{\mu^1, (\mu^2)^t}(Q_F; q, e^t)}{\mathcal{K}_{(0)(0)}(Q_F; q, e^{-t})\mathcal{K}_{(0)(0)}(Q_F; q, e^t)}. \end{aligned}$$

Now we can apply Theorem 5.1 and (11). $\square$

By the second product expression in Theorem 5.1 one can also express

$$\frac{\mathcal{Z}_{\mathbb{F}_m}(q, Q_B, Q_F, e^{-t})}{\mathcal{K}_{(0)(0)}(Q_F; q, e^{-t})\mathcal{K}_{(0)(0)}(Q_F; q, e^t)}$$

as a sum of infinite products. This expression generalizes Nekrasov's partition function.

We propose the deformed partition function to be:

$$\begin{aligned} & \mathcal{Z}_{\mathbb{F}_m}(Q_B, Q_F; q, e^{-t}) \\ &= \sum_{\mu^1, 2, \nu^1, 2} (-1)^{m(|\mu^1|+|\mu^2|)} q^{-m\kappa_{\mu^1}} q^{m\kappa_{\mu^2}/2} Q_B^{|\mu^1|} (Q_B Q_F^m)^{|\mu^2|} \\ & \quad \cdot W_{\mu^1 \nu^1}(q, e^{-t}) Q_F^{|\nu^1|} W_{\nu^1 \mu^2}(q, e^{-t}) \cdot W_{\mu^2 \nu^2}(q, e^{-t}) Q_F^{|\nu^2|} W_{\nu^2 \mu^1}(q, e^{-t}). \end{aligned}$$

We have checked in some low degree cases that this to satisfy the following deformed version of Gopakumar-Vafa integrality:

$$(101) \quad F = \sum_{\Sigma \in H_2(X) - \{0\}} \sum_{g \geq 0} \sum_{k \geq 1} \frac{1}{k} (-1)^{g-1} n_{\Sigma}^g (e^{-kt}) (q^{\frac{k}{2}} - q^{-\frac{k}{2}})^{2g-2} Q^{k\Sigma},$$

where $n_{\Sigma}^g(e^{-t})$ is a polynomial in e^{-t} with integral coefficients. For example, for $\mathbb{F}_0$, we have

$$\begin{aligned} n_B^g(e^{-t}) &= n_F^g = -2(1 - e^{-t})^2 \delta_{g,0}, \\ n_{B^2}^g &= n_{F^2}^g = 0, \quad n_{B+F}^g = -4(1 - e^{-t})^3, \\ n_{B^3}^g &= n_{F^3}^g = 0, \quad n_{2B+F}^g = n_{B+2F}^g = -(6 - 2e^{-t})(1 - e^{-t})^3 \delta_{g,0}. \end{aligned}$$

For $\mathbb{F}_1$ we have

$$\begin{aligned} n_B^g &= (1 - e^{-t})^2 \delta_{g,0}, & n_F^g &= -2(1 - e^{-t})^2 \delta_{g,0}, \\ n_{2B}^g &= n_{2F}^g = 0, & n_{B+F}^g &= (1 - e^{-t})^3 (3 + e^{-t}) \delta_{g,0}, \\ n_{3B}^0 &= -2e^{-3t} (1 - e^{-t})^3 \delta_{g,0}, & n_{B^3}^1 &= -e^{-4t}, \\ n_{2B+F}^g &= (1 - e^{-t})^2 (2e^{-2t} + 2e^{-3t}) \delta_{g,0}, \\ n_{B+2F}^g &= (1 - e^{-t})^2 (5 - 6e^{-t} - e^{-2t}) \delta_{g,0}. \end{aligned}$$

For $\mathbb{F}_2$ we have

$$\begin{aligned} n_B^g &= -(1 - e^{-t})^2 \delta_{g,0}, & n_F^g &= -2(1 - e^{-t})^2 \delta_{g,0}, \\ n_{2B}^0 &= 2e^{-t} (1 - e^{-t})^3, & n_{2B}^0 &= e^{-t} (1 - e^{-t})^2, \\ n_{B+F}^g &= -2(1 - e^{-t})^3 \delta_{g,0}, & n_{2F}^g &= 0. \end{aligned}$$

In general, one can consider the deformed topological string partition functions for toric local Calabi-Yau 3-folds in the same fashion and study the deformed Gopakumar-Vafa invariants. For example, for $\mathbb{P}^2$ we have

$$\begin{aligned} n_1^g(e^{-t}) &= 3(1 - e^{-t})^2 \delta_{g,0}, \\ n_2^g(e^{-t}) &= -3(1 - e^{-t})^2 (2 - e^{-t}) \delta_{g,0}, \\ n_3^0(e^{-t}) &= (27 - 15e^{-t})(1 - e^{-t})^3, & n_3^1 &= (-10 + 6e^{-t})(1 - e^{-t})^2, & n_3^g &= 0 (g > 1), \\ n_4^0 &= -(1 - e^{-t})^2 (192 - 474e^{-t} + 390e^{-2t} - 114e^{-3t} + 6e^{-4t}), \\ n_4^1 &= (1 - e^{-t})^2 (231 - 402e^{-t} + 201e^{-2t} - 24e^{-3t}), \\ n_4^2 &= -(1 - e^{-t})^2 (102 - 120e^{-t} + 30e^{-2t}), \\ n_4^3 &= (1 - e^{-t})^2 (15 - 12e^{-t}), \\ n_4^g &= 0, \quad g > 3. \end{aligned}$$

We notice the following positivity of n_d^g for these example. Define

$$P_d^g(x) = (-1)^{g+d-1} n_d^g(-x).$$

Then the coefficients of $P_d^g(x)$ are nonnegative. We conjecture it is true in general.

Here we want to offer some speculations on the geometric meaning of extra variable in the partition function. For a toric Fano surface X , the mathematical definition of its Gromov-Witten invariants are defined as follows:

$$\int_{[\overline{\mathcal{M}}_{g,0}(X,d)]^{\text{virt}}} e(K_X^{g,d}),$$

where $d \in H_2(X, \mathbb{Z})$, $\overline{\mathcal{M}}_{g,0}(X, d)$ is the moduli space of genus g stable maps of degree d to X , $K_X^{g,d}$ is the vector bundle on $\overline{\mathcal{M}}_{g,0}(X, d)$ whose fiber at a map $f : \Sigma \rightarrow X$ is given by $H^1(\Sigma, f^* K_X)$. Since $\overline{\mathcal{M}}_{g,0}(X, d)$ is compact, one can do the calculations in the equivariant setting:

$$\int_{[\overline{\mathcal{M}}_{g,0}(X,d)]_{T^2}^{\text{virt}}} e_{T^2}(K_X^{g,d}),$$

and the result is a constant. Here we use the T^2 -action that defines the toric structure. However, it is tempting also to do the calculation equivariantly directly

on K_X . For this purpose consider an extra circle action on the fiber of the K_X by multiplication and consider

$$\int_{[\mathcal{M}_{g,0}(K_X,d)]_{T^3}^{\text{virt}}} 1.$$

One can take the result obtained by formally applying the localization as definition. This might explain the extra parameter in the deformed partition functions.

6.5. The $SU(N)$ ($N > 2$) cases. The undeformed topological string partition function is given by the topological vertex as follows [10]:

$$(102) \quad \begin{aligned} Z_{A_{N-1}}^{(m)} &= \sum_{\mu^1, \dots, \mu^N} \prod_{i=1}^N (Q_{b_i}^{|\mu^i|} (-1)^{(N+m-2i)|\mu^i|} q^{(N+2m-2i+1)\kappa_{\mu^i}}) \\ &\cdot \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N (\mathcal{W}_{\nu^{k-1} \mu^k (\nu^k)^t} (q) q^{\kappa_{\nu^k/2}} Q_k^{|\nu^k|}) \\ &\cdot \sum_{\eta^1, \dots, \eta^{N-1}} \prod_{k=1}^N (\mathcal{W}_{\eta^{k-1} (\mu^{N+1-k})^t (\eta^k)^t} (q) q^{\kappa_{\eta^k/2}} Q_{N+1-k}^{|\eta^k|}), \end{aligned}$$

where $\nu^0 = \nu^N = \eta^0 = \eta^N = (0)$, $Q_N = 1$. This suggests the following deformed partition function:

$$(103) \quad \begin{aligned} \mathcal{Z}_{A_{N-1}}^{(m)} &= \sum_{\mu^1, \dots, \mu^N} \prod_{i=1}^N (Q_{b_i}^{|\mu^i|} (-1)^{(N+m-2i)|\mu^i|} q^{(N+2m-2i+1)\kappa_{\mu^i}}) \\ &\cdot \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N (W_{\nu^{k-1} \mu^k (\nu^k)^t} (q, e^{-t}) q^{\kappa_{\nu^k/2}} Q_k^{|\nu^k|}) \\ &\cdot \sum_{\eta^1, \dots, \eta^{N-1}} \prod_{k=1}^N (W_{\eta^{k-1} (\mu^{N+1-k})^t (\eta^k)^t} (q, e^{-t}) q^{\kappa_{\eta^k/2}} Q_{N+1-k}^{|\eta^k|}), \end{aligned}$$

where $\nu^0 = \nu^N = \eta^0 = \eta^N = (0)$, $Q_N = 1$. We expect this to satisfy the deformed Gopakumar-Vafa integrality.

On the other hand, one can use the following symmetry of the topological vertex (see e.g. [23]):

$$\mathcal{W}_{(\mu^1)^t (\mu^2)^t (\mu^3)^t} (q) = q^{-(\kappa_{\mu^1} + \kappa_{\mu^2} + \kappa_{\mu^3})/2} \mathcal{W}_{\mu^3, \mu^2, \mu^1} (q)$$

to rewrite the last line in (102). This yields the following expression [10]:

$$(104) \quad \begin{aligned} Z_{A_{N-1}}^{(m)} &= \sum_{\mu^1, \dots, \mu^N} \prod_{i=1}^N (Q_{b_i}^{|\mu^i|} (-1)^{(N+m)|\mu^i|} q^{(N+2m-2i)\kappa_{\mu^i}/2}) \\ &\cdot \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N (\mathcal{W}_{\nu^{k-1} \mu^k (\nu^k)^t} (q) q^{\kappa_{\nu^k/2}} Q_k^{|\nu^k|}) \\ &\cdot \sum_{\eta^1, \dots, \eta^{N-1}} \prod_{k=1}^N (\mathcal{W}_{\eta^{k-1} \mu^k (\eta^k)^t} (q) q^{\kappa_{\eta^k/2}} Q_k^{|\eta^k|}). \end{aligned}$$

This suggests the following partition function:

$$\begin{aligned}
 \tilde{Z}_{A_{N-1}}^{(m)} &= \sum_{\mu^1, \dots, \mu^N} \prod_{i=1}^N (Q_{b_i}^{|\mu^i|} (-1)^{(N+m)|\mu^i|} q^{(N+2m-2i)\kappa_{\mu^i}/2}) \\
 (105) \quad &\cdot \sum_{\nu^1, \dots, \nu^{N-1}} \prod_{k=1}^N (W_{\nu^{k-1} \mu^k (\nu^k)^t} (q, e^{-t}) q^{\kappa_{\nu^k}/2} Q_k^{|\nu^k|}) \\
 &\cdot \sum_{\eta^1, \dots, \eta^{N-1}} \prod_{k=1}^N (W_{\eta^{k-1} \mu^k (\eta^k)^t} (q, e^t) q^{\kappa_{\eta^k}/2} Q_k^{|\eta^k|}).
 \end{aligned}$$

In other words,

$$\begin{aligned}
 \tilde{Z}_{A_{N-1}}^{(m)} &= \sum_{\mu^1, \dots, \mu^N} \prod_{i=1}^N (Q_{b_i}^{|\mu^i|} (-1)^{(N+m)|\mu^i|} q^{(N+2m-2i)\kappa_{\mu^i}/2}) \\
 &\cdot \tilde{\mathcal{K}}_{\mu^1, \dots, \mu^N} (Q_1, \dots, Q_{N-1}; q, e^{-t}) \cdot \tilde{\mathcal{K}}_{\mu^1, \dots, \mu^N} (Q_1, \dots, Q_{N-1}; q, e^t)
 \end{aligned}$$

As in the $N = 2$ case one can try to relate it to the generating series of

$$\chi(M(N, k), \Lambda_{-e^{-t}}(\mathbb{V}_k \otimes \mathbb{W}_k^*) \otimes \Lambda_{-e^{-t}}(\mathbb{V}_k^* \otimes \mathbb{W}_k) \otimes E_{N,k}^m)(e_1, \dots, e_N, q, q^{-1}).$$

Acknowledgements. This research is partially supported by research grants from NSFC and Tsinghua University. The final stage of its preparation was carried out in the Center of Mathematical Sciences, Zhejiang University. The author thanks Professor Kefeng Liu for his interest in this work.

REFERENCES

- [1] M. Aganagic, A. Klemm, M. Marino, C. Vafa, *The topological vertex*, hep-th/0305132.
- [2] M. Aganagic, M. Mariño, C. Vafa, *All loop topological string amplitudes from Chern-Simons theory*, hep-th/0206255.
- [3] G.E. Andrews, *The theory of partitions*. Encyclopedia of Mathematics and its Applications, Vol. 2. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976.
- [4] H. Awata, H. Kanno, *Instanton counting, Macdonald function and the moduli space of D-branes*, hep-th/0502061.
- [5] T. Eguchi, H. Kanno, *Topological Strings and Nekrasovs formulas*, hep-th/0310235.
- [6] T. Eguchi, H. Kanno, *Geometric transitions, Chern-Simons gauge theory and Veneziano type amplitudes*, hep-th/0312234.
- [7] T. Holowood, A. Iqbal, C. Vafa, *Matrix models, geometric engineering and elliptic genera*, hep-th/0310272.
- [8] A. Iqbal, *All genus topological string amplitudes and 5-brane webs as Feynman diagrams*, hep-th/0207114.
- [9] A. Iqbal, A.-K. Kashani-Poor, *SU(N) geometries and topological string amplitudes*, hep-th/0306032.
- [10] A. Iqbal, A.-K. Kashani-Poor, *Instanton counting and Chern-Simons theory*, hep-th/0212279.
- [11] J. Li, C.-C. Liu, K. Liu, J. Zhou, *A mathematical theory of the topological vertex*, math.AG/0408426.
- [12] J. Li, K. Liu, J. Zhou, *Topological string partition functions as equivariant indices*, math.AG/0412089.
- [13] I.G. MacDonald, *Symmetric functions and Hall polynomials*, 2nd edition. Clarendon Press, 1995.
- [14] M. Mariño, C. Vafa, *Framed knots at large N*, Orbifolds in mathematics and physics (Madison, WI, 2001), 185–204, Contemp. Math., 310, Amer. Math. Soc., Providence, RI, 2002.

- [15] h.R. Moron, S.G. Lukac, *The Homfly plonomial of the decorated Hopf link*, J. Knot Theory Ramif. 12 (2003), 395-416, math.GT/0108011.
- [16] H. Nakajima, K. Yoshioka, *Instanton counting on blowup, I*, math.AG/0306198.
- [17] N. A. Nekrasov, *Trieste lectures on solitons in nocommutative gauge theory*, hep-th/0011095.
- [18] N. A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, hep-th/0206161.
- [19] N. Nekrasov, A. Okounkov, *Seiberg-Witten theory and random partitions*, hep-th/0306238.
- [20] A. Okounkov, N. Reshtikhin, C. Vafa, *Quantum Calabi-Yau and classical crystals*, hep-th/0309208.
- [21] J. Zhou, *A conjecture on Hodge integrals*, math.AG/0310282.
- [22] J. Zhou, *Localizations on moduli Spaces and free field realizations of Feynman rules*, math.AG/0310283.
- [23] J. Zhou, *Curve counting and instanton counting*, to appear in Adv. Theoret. Math. Phys., math.AG/0311237.

CENTER OF MATHATICAL SCIENCES, ZHEJIANG UNIVERSITY, AND DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, CHINA

E-mail address: jzhou@math.tsinghua.edu.cn